



**Australian Government**  
**Department of Health**

# COMMUNICABLE DISEASES INTELLIGENCE

2020      Volume 44  
<https://doi.org/10.33321/cdi.2020.44.62>

## **Australian Meningococcal Surveillance Programme annual report, 2019**

Monica M Lahra and Tiffany R Hogan for the National Neisseria Network

# Communicable Diseases Intelligence

ISSN: 2209-6051 Online

This journal is indexed by Index Medicus and Medline.

Creative Commons Licence - Attribution-NonCommercial-NoDerivatives CC BY-NC-ND

© 2020 Commonwealth of Australia as represented by the Department of Health

This publication is licensed under a Creative Commons Attribution-Non-Commercial NoDerivatives 4.0 International Licence from <https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode> (Licence). You must read and understand the Licence before using any material from this publication.

## Restrictions

The Licence does not cover, and there is no permission given for, use of any of the following material found in this publication (if any):

- the Commonwealth Coat of Arms (by way of information, the terms under which the Coat of Arms may be used can be found at [www.itsanhonour.gov.au](http://www.itsanhonour.gov.au));
- any logos (including the Department of Health's logo) and trademarks;
- any photographs and images;
- any signatures; and
- any material belonging to third parties.

## Disclaimer

Opinions expressed in Communicable Diseases Intelligence are those of the authors and not necessarily those of the Australian Government Department of Health or the Communicable Diseases Network Australia. Data may be subject to revision.

## Enquiries

Enquiries regarding any other use of this publication should be addressed to the Communication Branch, Department of Health, GPO Box 9848, Canberra ACT 2601, or via e-mail to: [copyright@health.gov.au](mailto:copyright@health.gov.au)

## Communicable Diseases Network Australia

Communicable Diseases Intelligence contributes to the work of the Communicable Diseases Network Australia.  
<http://www.health.gov.au/cdna>



Communicable Diseases Intelligence (CDI) is a peer-reviewed scientific journal published by the Office of Health Protection, Department of Health. The journal aims to disseminate information on the epidemiology, surveillance, prevention and control of communicable diseases of relevance to Australia.

## Editor

Tanja Farmer

## Deputy Editor

Simon Petrie

## Design and Production

Kasra Yousefi

## Editorial Advisory Board

David Durrheim,  
Mark Ferson, John Kaldor,  
Martyn Kirk and Linda Selvey

## Website

<http://www.health.gov.au/cdi>

## Contacts

Communicable Diseases Intelligence is produced by:  
Health Protection Policy Branch  
Office of Health Protection  
Australian Government  
Department of Health  
GPO Box 9848, (MDP 6)  
CANBERRA ACT 2601

## Email:

[cdi.editor@health.gov.au](mailto:cdi.editor@health.gov.au)

## Submit an Article

You are invited to submit your next communicable disease related article to the Communicable Diseases Intelligence (CDI) for consideration. More information regarding CDI can be found at:  
<http://health.gov.au/cdi>.

Further enquiries should be directed to:  
[cdi.editor@health.gov.au](mailto:cdi.editor@health.gov.au).

# Australian Meningococcal Surveillance Programme annual report, 2019

Monica M Lahra and Tiffany R Hogan for the National Neisseria Network

## Abstract

Invasive meningococcal disease (IMD) is a notifiable disease in Australia, and both probable and laboratory-confirmed cases of IMD are reported to the National Notifiable Diseases Surveillance System (NNDSS). In 2019, there were 206 notifications of IMD. Of these, 202 were laboratory-confirmed cases analysed by the reference laboratories of the Australian National Neisseria Network (NNN). Of the 202 laboratory-confirmed cases of IMD, 167 were confirmed by bacterial culture and 35 by nucleic acid amplification testing, and all had the serogroup determined. Fine typing was available on 146 samples (146/202, 72%). *Neisseria meningitidis* serogroup B (MenB) infections accounted for 50.0% (101/202); MenW for 26.2% (53/202); MenY for 20.8% (42/202) and MenC for 3.0% of cases (6/202). Of the MenW cases, 88% were PorA antigen type P1.5,2, and 65% of these (24/37) were sequence type 11, the hypervirulent strain reported in recent outbreaks in Australia and overseas.

The primary peaks of IMD notifications in Australia in 2019 were observed in infants less than 1 year of age (36/202, 18%) and in adults aged 65 years or older (39/202, 19%). MenB infections predominated in those aged less than 5 years and those aged 15–19 years, whereas MenW and MenY infections predominated in those aged 45 years or more.

All 167 IMD isolates were tested for antimicrobial susceptibility. One isolate out of these 167 (0.6%) was resistant to penicillin with an MIC  $\geq$  1mg/L; 154/167 isolates (92%) had decreased susceptibility to penicillin. All isolates were susceptible to ceftriaxone and ciprofloxacin, and one isolate was resistant to rifampicin.

Keywords: antibiotic resistance; disease surveillance; meningococcal disease; *Neisseria meningitidis*

## Introduction

Australia's National Neisseria Network (NNN) was established in 1979 as a collaborative network of reference laboratories in each state and territory which contribute to the laboratory surveillance of the pathogenic *Neisseria*: *N. meningitidis* and *N. gonorrhoeae*. The NNN has coordinated laboratory data from cases of invasive meningococcal disease (IMD) for the Australian Meningococcal Surveillance Programme (AMSP) since 1994, and is supported by the Australian Government Department of Health and the jurisdictions.<sup>1</sup> The NNN laboratories supply phenotypic and genotypic data to sup-

plement the notification data from the National Notifiable Diseases Surveillance System (NNDSS), which includes cases of probable and laboratory-confirmed IMD.

Notifications of IMD in Australia peaked in 2002 at 3.5 cases per 100,000,<sup>2</sup> with the majority of disease at that time caused by MenB and MenC. In 2003, the introduction of the conjugate serogroup C meningococcal vaccine to the National Immunisation Program was followed by significant and sustained reductions, in the 10 years that followed, of both the number of serogroup C IMD cases and the overall notifications of IMD, to a rate of only 0.6 cases per 100,000 in 2013.<sup>3,4</sup>

After 2013, however, there followed increases both in IMD notifications and in the proportion of IMD caused by the MenW and MenY serogroups in Australia. The IMD notification rate increased to 1.5 cases per 100,000 in 2017,<sup>2</sup> when MenACWY immunisation programmes were implemented across jurisdictions in targeted age groups. This was followed by a change in the national immunisation programme in 2018 substituting monovalent MenC vaccine with the 4-valent MenACWY vaccine. In 2018 IMD notifications declined to 1.1 per 100,000, then to 0.8 per 100,000 in 2019.

IMD is a rare disease in Australia, but one of public health concern. Continued monitoring of phenotypic and genotypic features of IMD strains is critical to plan and inform clinical management of cases, case clusters and outbreaks of IMD locally and nationally, and to inform and monitor public health interventions.

## Methods

### Case confirmation of invasive meningococcal disease

Case confirmation is based on isolation of *N. meningitidis*, or on a positive nucleic acid amplification testing (NAAT) from a normally sterile site, defined as laboratory-definitive evidence of IMD according to the national case definition.<sup>5</sup> Information regarding the site of infection, age and sex of the patients is collated by the NNN for the AMSP.

IMD cases are categorised on the basis of the site from which *N. meningitidis* was isolated, or from which meningococcal DNA was detected (blood, joint fluid, vitreous fluid). When *N. meningitidis* is detected from both blood and cerebrospinal fluid (CSF) from the same patient, the case is classified as one of meningitis.

## Phenotyping and genotyping of *Neisseria meningitidis*

Phenotyping is limited to the determination of the serogroup by detection of soluble polysaccharide antigens. Genotyping of both isolates and DNA extracts is performed by sequencing of products derived from amplification of the porin genes *porA* and *porB* and the outer membrane protein, *fetA*.

## Antibiotic susceptibility testing


Meningococcal isolates are tested to determine their minimum inhibitory concentration (MIC) values to antibiotics used for therapeutic and prophylactic purposes: ceftriaxone, ciprofloxacin, penicillin and rifampicin. This program has historically reported penicillin testing categories as: sensitive (MIC  $\leq$  0.03 mg/L); less sensitive (MIC 0.06–0.5 mg/L); and resistant (MIC  $\geq$  1 mg/L). However, to monitor across antimicrobial susceptibility testing methods, a distribution of penicillin MIC values is now reported.

## Results

In 2019, there were 202 laboratory-confirmed cases of IMD analysed by the NNN, and 206 cases notified to the NNDSS.<sup>2</sup> The numbers of notifications has continued to decline following the recent MenACWY vaccination interventions in Australia – from 380 in 2017 to 281 in 2018 and 206 in 2019. Laboratory data were available for 98% of notified cases of IMD in Australia in 2019 (Figure 1). In 2019, the peak incidence for IMD occurred in midwinter and early spring (1 July to 30 September) (Table 1).

In 2019 there was a decrease in notifications of IMD across jurisdictions compared to 2018. While New South Wales reported the highest number of cases (58 cases), this was a decrease from 70 cases in 2018. Queensland had the second highest number of notifications (45 cases), a decrease from 58 cases in 2018. The number of cases from each jurisdiction for 2019 is shown in Table 2.

**Figure 1: Number of invasive meningococcal disease cases reported to the National Notifiable Diseases Surveillance System compared with laboratory-confirmed data from the Australian Meningococcal Surveillance Programme, Australia, 1991–2019**



**Table 1: Laboratory-confirmed cases of invasive meningococcal disease, Australia 2019**

IMD serogroup 2019	1 January – 31 March	1 April – 30 June	1 July – 30 September	1 October – 31 December	Total
B	19	25	37	20	101
C	3	2	0	1	6
Y	4	8	21	9	42
W	11	13	19	10	53
<b>Total</b>	<b>37</b>	<b>48</b>	<b>77</b>	<b>40</b>	<b>202</b>

**Table 2: Number of laboratory-confirmed cases of invasive meningococcal disease, Australia, 2019, by state or territory and serogroup**

State/ territory	Serogroups				Total
	B	C	Y	W	
ACT	1	0	0	0	1
NSW	34	0	14	10	58
NT	1	0	0	6	7
Qld	18	0	18	9	45
SA	18	0	4	3	25
Tas	5	0	0	2	7
Vic	16	0	4	14	34
WA	8	6	2	9	25
<b>Australia</b>	<b>101</b>	<b>6</b>	<b>42</b>	<b>53</b>	<b>202</b>
	50.0	3.0	20.8	26.2	%

**Table 3: Laboratory-confirmed cases of invasive meningococcal disease, by age and serogroup; and the proportion of IMD attributable to MenB Australia, 2019**

Serogroup	Age group									Total
	<1	1–4	5–9	10–14	15–19	20–24	25–44	45–64	65+	
B	20	13	3	6	21	10	12	12	4	101
C	1	0	1	0	0	0	3	1	0	6
Y	4	0	0	1	3	9	0	9	16	42
W	11	1	2	0	2	2	9	7	19	53
<b>Total</b>	<b>36</b>	<b>14</b>	<b>6</b>	<b>7</b>	<b>26</b>	<b>21</b>	<b>24</b>	<b>29</b>	<b>39</b>	<b>202</b>
%B of within age group	55.6	92.9	50.0	85.7	80.8	47.6	50.0	41.4	10.3	

**Table 4: Number of laboratory-confirmed cases of invasive meningococcal disease, Australia, 2019, by specimen type and method of confirmation**

Specimen	Bacterial culture	Nucleic Acid Amplification Test	Total
Blood	150	14	164
CSF +/- Blood	13	20	33
Other specimens <sup>a</sup>	4	1	5
<b>Total</b>	<b>167</b>	<b>35</b>	<b>202</b>

a Other specimens: joint aspirates (3); ascitic fluid (1); and unspecified (1).

## Age distribution

The peak incidence of IMD in 2019 occurred in the extremes of age: in infants of less than one year (18% of cases, 36/202); and adults aged over 65 years or more, comprising 19% of IMD cases (38/202) in 2019 (Table 3). We note also that 25% of IMD occurred in children less than 5 years of age (50/202). Between 2003 and 2014, the proportion of IMD that occurred in children aged less than 5 years ranged from 28% to 36% of cases. Since 2015, in Australia the proportion of IMD in children less than 5 years of age has ranged from 21% to 27%.

## Samples from laboratory-confirmed cases

In 2019, diagnosis was confirmed by a positive bacterial culture in 83% of cases of IMD (167/202); for 17% (35/202) of cases, IMD was confirmed by NAAT testing alone (Table 4). There were 33 diagnoses of meningitis and 164 diagnoses of septicaemia (Table 4). Additionally, there were 5 IMD diagnoses confirmed from other sites comprising joint fluid aspirates (n = 3), ascitic fluid (n = 1), and an unspecified site (n = 1).

## Notifications and proportion of MenB, MenC, MenY and MenW

The serogroup was determined for all 202 laboratory-confirmed cases of IMD (Tables 2 and 3). In 2019, MenB accounted for 50% of all IMD in Australia. With respect to serogroup infections by age group, as shown in Table 3, in children aged 1–4 years (13/14, 92.9%) and in older chil-

dren and adolescents (10–19 years) (27/33, 81.8%) MenB predominated. MenB accounted for the largest proportion of disease in all age groups (range 41.4–92.9%) excepting 65 years and older where MenW and MenY were in the majority.

New South Wales reported the largest number of notifications of MenB IMD (n = 34), accounting for 59% of IMD in this state (34/58) in 2019. South Australia was again the state with the highest proportion of MenB, 72% of statewide IMD cases (18/25). In the years 2006–2012 the proportion of IMD cases caused by MenB was 84–88%, before falling in successive years: in 2013–2014 to 75–80%, in 2015 to 64%, and to 36% in 2016–2017. There was an increase in its attributable proportion in 2018 to 44% and again in 2019 to 50% of IMD cases reported (Figure 2).

There were 6 notifications of IMD caused by MenC in 2019; this is higher than the number reported in 2018 (n = 4), but similar to annual national case numbers for the period 2014–2016. All MenC IMD cases in 2019 were reported from Western Australia (Table 3, Figure 3).

The rise in IMD notifications in Australia since 2014 is due to a rise in the number of infections of both MenW and MenY, as shown in Figure 1 and Figure 2. Prior to 2015 the proportion of cases of IMD caused by MenW was low, ranging from 1.1–4.8% in 1997–2012, then increasing to 8.4–9.7% in 2013–2014. In 2015 it was 21%, in 2016 it was 44%, in 2017–2018 it was 36–38%, as shown in Figure 2. In 2019 there were 53/202 MenW infections, accounting for 26% of IMD.

Figure 2: Proportion of serogroups of laboratory-confirmed invasive meningococcal disease, Australia, by year




Figure 3: Number of serogroup B, C, Y and W cases of laboratory-confirmed invasive meningococcal disease, Australia, 2019, by age




Figure 4: Number of *porA* genotypes for MenB in laboratory-confirmed cases of invasive meningococcal disease Australia, 2019




Figure 5: Number of *porA* genotypes for MenW in laboratory-confirmed cases of invasive meningococcal disease Australia, 2019





Figure 6: Number of *porA* genotypes for MenY in laboratory-confirmed cases of invasive meningococcal disease Australia, 2019



**Table 5: Distribution of *porA* genotype laboratory-confirmed cases of invasive meningococcal disease, Australia, 2019, by state or territory**

<i>porA</i> genotype	Number per serogroup per state / territory							
	NSW	Qld	Vic	SA	WA	ACT	Tas	NT
P1.5,2	3W	9W	11W	2W	7W, 5C	0	2W	3W
P1.5,2-67	0	0	0	0	1W	0	0	0
P1.5,10-1	0	1Y	0	0	0	0	0	0
P1.5-1,2-2	0	0	2W	0	0	0	0	0
P1.5-1,5-1	0	1Y	0	0	0	0	0	0
P1.5-1,10-1	1Y	14Y	3Y, 1B	2Y	2Y	0	0	0
P1.5-1,10-4	0	2Y	1Y, 1W	0	0	0	0	0
P1.5-2,10-29	1Y	0	0	0	0	0	0	0
P1.7,16-26	3B	3B	4B	1B	1B	0	0	0
P1.7,16-29	0	0	0	0	1C	0	0	0
P1.7, 16-44	1B	0	0	0	0	0	0	0
P1.7-2,4	3B	1B	3B	8B	1B	0	0	0
P1.7-2,5	0	0	0	1B	0	0	0	0
P1.7-2,6	0	0	0	1B	0	0	0	0
P1.7-2,16-26	1B	0	0	1B	0	0	0	0
P1.7-4,1	0	1B	0	0	0	0	0	0
P1.7-12,14	0	2B	0	0	0	0	0	0
P1.12-1,16	2B	0	0	0	0	0	0	0
P1.12-6,16-169	0	1B	0	0	0	0	0	0
P1.18-1,16	1W	0	0	0	0	0	0	0
P1.18-1,34	0	1B	3B	0	2B	0	1B	0
P1.19,15	0	0	1B	0	0	0	1B	0
P1.19-2,13-1	0	1B	0	1B	0	0	0	0
P1.22,9	0	0	1B	1B	1B	0	0	0
P1.22,14	1B	7B	3B	0	0	1B	1B	0
P1.22,14-3	0	1B	0	0	0	0	0	0
P1.22,14-6	0	0	0	0	1B	0	0	0
P1.22-1, 14	0	0	0	1B	0	0	0	0

**Table 6: Laboratory-confirmed cases of MenW invasive meningococcal disease, Australia, 2019, by sequence type (ST)**

Sequence type	MenW genotype					Total
	P1.5,2	P1.5,2-67	P1.5-1,2-2	P1.5-1,10-4	P1.18-1,16	
ST-11	24	1	0	1	0	<b>26</b>
ST-22	0	0	2	0	0	<b>2</b>
ST574	0	0	0	0	0	<b>0</b>
ST1158	0	0	0	0	0	<b>0</b>
ST1287	0	0	0	0	0	<b>0</b>
ST1655	0	0	0	0	0	<b>0</b>
ST2780	0	0	0	0	0	<b>0</b>
ST12351	0	0	0	0	0	<b>0</b>
ST13135	0	0	0	0	0	<b>0</b>
Not determined	13	0	0	0	1	<b>14</b>
<b>Total</b>	<b>37</b>	<b>1</b>	<b>2</b>	<b>1</b>	<b>1</b>	<b>42</b>

IMD caused by MenW was seen in all age groups in 2019 except those aged between 10 to 14 years. We note that this age group has historically had low numbers of IMD cases; in 2019, only 7 cases (6 MenB and 1 MenY) were reported. For those aged less than 5 years, the number and proportion of IMD cases caused by MenW (12 cases, 24%) was lower than that reported in 2018. For those aged over 65 years, MenW was the predominant cause of IMD.

A similar pattern and trend was seen in the number and proportion of IMD caused by MenY in 2019 (n = 42, 21% of total IMD). Prior to 2015, the proportion of cases of IMD caused by MenY ranged from 1.3–4.6% in the period 1997–2010 to 6.2–10.5% in 2011–2014. In 2015 the proportion rose to 12.6%, increasing to 16.5% in 2016 and again to 20% in 2017. The proportion was lower in 2018, at 15.8% (Figure 2).

Of the 42 cases of IMD caused by MenY in 2019, Queensland reported the largest number (18/45 statewide notifications in 2019). MenY was reported in all jurisdictions except the Australian Capital Territory, Northern Territory and Tasmania (Table 2). MenY IMD was reported in all age groups in 2019 except in

those aged between 1 to 9 years and 25 to 44 years, with the highest number and proportion seen in those aged 45 or more years (25/68 of the total IMD notifications for this age group), as in previous years (Table 3, Figure 3).

### Genotyping

In 2019, genotyping was possible for 72% of IMD notifications (146/202). Results are shown in Figures 4–6 and Table 5. There were 70 MenB cases typed: the predominant *porA* type in 2019 continues to be P1.7-2,4 (16/70, 23%). Other prevalent MenB genotypes were P1.22,14 (13 cases, 19%), and P1.7,16-26 (12 cases, 17%), similar to previous years (Table 5). All MenC IMD in 2019 were from Western Australia, and investigation revealed that 5/6 were type P1.5,2 (Table 5). For serogroup W IMD, the predominant genotype remains P1.5,2 (37 cases, 88%) (Figures 4–6 and Table 5). Of these, 24 cases (65%) were clonal complex 11, the same strain type as the hyper-virulent serogroup W strain also reported in the UK and South America since 2009 (Table 6).<sup>6,7</sup> For MenY IMD, the predominant genotype was P1.5-1,10-1 (22 cases, 79%), as has been reported since 2014, when the increase in serogroup Y IMD was first noted in Australia (Figure 6).

## Antibiotic susceptibility testing

Isolates of *N meningitidis* are tested against both treatment (ceftriaxone and penicillin) and clearance antibiotics (rifampicin and ciprofloxacin). In 2019, 167/202 of laboratory-confirmed IMD had *Neisseria meningitidis* cultured, and it was therefore possible for antimicrobial susceptibility testing (AST) to be performed. All 167 *N. meningitidis* isolates had AST performed by the NNN.

Regarding antimicrobial susceptibility to the treatment antibiotics, ceftriaxone susceptibility testing was performed on all 167 isolates in 2019, and all isolates were susceptible. With regards to penicillin, the distribution of penicillin MIC values is shown in Table 7. There were 12/167 strains (7.2%) fully susceptible to penicillin ( $\text{MIC} \leq 0.03 \text{ mg/L}$ ).

There were 154/167 isolates that were categorised as less susceptible to penicillin (MIC value 0.064–0.5 mg/L). Of these, 34 had a penicillin MIC of 0.5mg/L, which is close to the MIC breakpoint for resistance; 26/34 (76%) with this penicillin MIC were Men W. In recent years MenW has demonstrated higher penicillin MIC values and higher proportions of resistance. In 2019, one of 167 IMD isolates (0.6%) was resistant to penicillin ( $\text{MIC} \geq 1 \text{ mg/L}$ ): this isolate was a MenW IMD, diagnosed in Victoria. The proportion of IMD isolates resistant to penicillin was lower than reported in 2018 (1.4%) and continues to decline from 2016 (5.8%) and 2017 (5.1%) as shown in Figure 7. Regarding the clearance antibiotics for IMD, all 167 isolates were susceptible to ciprofloxacin; one MenW isolate from Victoria was resistant to rifampicin with a MIC value  $\geq 32 \text{ mg/L}$ .

## Discussion


In 2019, 98% of IMD notifications (202/206) were laboratory confirmed.<sup>3</sup> Overall there was a 27% decrease in notifications of IMD in Australia in 2019 compared to 2018. Following the increases in MenW and MenY disease in Australia during 2016–2017, there were jurisdictional, time-limited MenACWY vaccination programs for target age groups in 2017 and 2018.<sup>8</sup> From 1 July 2018, a change to the National Immunisation Program (NIP) was implemented, where MenC vaccine at 12 months of age was replaced with a quadrivalent ACWY vaccine. This change in the NIP in 2018 has been followed by a decrease in both notifications and proportion of MenW and MenY disease, and also an increase in the proportion of disease caused by MenB from 36% in 2016–2017 to 50% in 2019. We note that, following the introduction of the MenC vaccine in 2003 in Australia, the proportion of MenB IMD was 84–88% in the years 2006–2012 then declining, along with disease notifications until the emergence of MenW and MenY disease since 2014 in Australia.<sup>8</sup> In 2019 MenB accounted for 66% of IMD (33/50) notifications in those less than 5 years of age; 81% of IMD (21/26) in 15–19 year olds; 48% of IMD (10/21) in 20–24 year olds; 50% of IMD (12/24) in 25–44 year olds; and 41% of IMD (12/29) in 45–64 year olds. A recombinant multi-component meningococcal B vaccine has been available in Australia since 2014,<sup>9</sup> however this vaccine is not currently on the NIP.

In 2019, with regards to serogroup infections by age group, as shown in Table 3, MenB predominated in children aged 1–4 years of age (13/14, 92.9%) and in older children and adolescents (10–19 years) (27/33, 81.8%). MenB accounted for

**Table 7: Penicillin MIC distribution of laboratory-confirmed invasive meningococcal disease isolates, Australia, 2019**

Penicillin MIC distribution									
MIC mg/L	$\leq 0.032$	0.064	0.125	0.25	0.5	1	2	$\geq 4$	Total
Number	12	35	25	60	34	1	0	0	167
%	7.2	21.0	15.0	35.9	20.4	0.6	0	0	100

Figure 7: Proportion of penicillin-susceptible, less susceptible and resistant invasive meningococcal disease isolates, Australia, by year



the greatest proportion of IMD in all age groups (range 41.4–92.9%) excepting 65 years and older where MenW and MenY were predominant.

MenW caused 26.1% of IMD (53/202) in Australia, a decrease from 36% in 2018. MenW IMD was reported in all age groups, except those between 10 to 14 years. MenW infections accounted for 30.6% of IMD in babies less than one year of age (11/36 notifications), and 49% of IMD notifications in those aged over 65 years (19/39 notifications). MenY IMD in 2019 (n = 42/202) accounted for 20.8% of IMD nationally; it was reported in babies less than one year of age (4/42, 9.5%); in 15–19 year olds (3/26, 11.5%); in 20–24 year olds (9/21, 43%); in 45–64 year olds (9/29, 31%); and in those aged 65 and older (16/39, 41%).

The primary peaks of IMD notifications were observed in infants less than 1 year of age (36/202, 18%) and in adults aged 65 years or older (39/202, 19%). In infants, IMD was predominately caused by MenB (20/36) and MenW (11/36), whereas in those aged 65 and older, IMD was attributable to MenW and MenY predominantly (35/39 notifications amongst those aged 65 and older were MenW or MenY). Secondary disease peaks were observed in adolescents 15–19 years of age, due primarily to MenB (81%).

With regards to the prevailing IMD serogroups and genotypes, the number of IMD cases caused by MenB in 2019 was less than in 2018, but the proportion was higher (101/202, 50% in 2019 compared with 123/278, 44% in 2018). New South Wales reported the largest number of MenB notifications. However, South Australia continues to report the highest proportion of MenB. The predominant genotypes of Men B in Australia continue to be P1.7-2,4 (16/70), followed by genotype P1.22,14 (13/70) and genotype P1.7,16-26 (12/70).

The predominant circulating strain of MenW continues to be genotype P1.5,2 sequence type (ST)-11. This MenW strain emerged in the United Kingdom and South America in 2009; it has spread to account for 25% of IMD in the

UK in 2014/15 and 59% of all cases in Chile in 2012.<sup>6,7</sup> MenW ST11 is hypervirulent and associated with atypical clinical presentations, more severe disease, and a higher case fatality rate.<sup>7</sup> The initial increase in MenW overseas, and in Australia, was seen in older adults; it was subsequently reported in all age groups, particularly in adolescents and infants.<sup>10</sup>

The predominant MenY genotype since 2014 continues to be P1.5-1,10-1; previously, MenY genotype distribution was more heterogeneous. The emergence of MenY IMD has also been reported recently in Europe.<sup>11</sup> The phenotypic and genotypic characterization of the MenY isolates is ongoing by the NNN.

Antimicrobial susceptibility testing of IMD isolates in 2019 showed that penicillin resistance has reduced compared with recent years (one isolate, 0.6%). The incidence of penicillin resistance in *N. meningitidis* in Australia was less than 1% annually among IMD isolates tested in 1996–2014, rising to 3.4% in 2015 and 5.8% in 2016; it has fallen from 5.1% in 2017 through 1.4% in 2018. However the proportion of IMD isolates less sensitive to penicillin has been increasing from 62–75% in 1996–2006; 67–79% in 2007–2009; 78–88% in 2010–2015; to 90–94% in 2016–2019. The majority of penicillin-resistant meningococcal isolates are MenW. All IMD isolates tested in 2019 were susceptible to ceftriaxone and ciprofloxacin, while one isolate was resistant to rifampicin.

There has been a year-on-year decrease in IMD notifications from 1.5 cases per 100,000 in 2017; to 1.1 per 100,000 in 2018; to 0.8 cases per 100,000 in 2019. There has also been a decrease in the proportion of isolates attributable to MenW and MenY following changes in the National Immunisation Programme. The NNN is continuing to lead further investigations with the Commonwealth Department of Health and is closely monitoring the phenotypic and genotypic features of *N. meningitidis* causing IMD in Australia. Additional investigations including whole genome sequencing are in place to enhance IMD surveillance. The AMSP data

are used for informing treatment guidelines and disease prevention strategies; and to monitor the effect of interventions.

## Acknowledgements

Meningococcal isolates were received in the reference centres from many laboratories throughout Australia. The considerable time and effort involved in forwarding these isolates is recognised and these efforts are greatly appreciated. These data could not have been provided without this assistance and the help of clinical colleagues and public health personnel. The Australian Government Department of Health provided funding for the National Neisseria Network.

Members of the AMSP in 2019, to whom isolates and samples should be referred, and enquiries directed, are listed below.

### Australian Capital Territory

P Collignon, S Bradbury  
Microbiology Department, The Canberra  
Hospital, Gilmore Crescent, Garran ACT 2605  
Telephone: +61 2 6244 2510  
Email: [peter.collignon@act.gov.au](mailto:peter.collignon@act.gov.au)

### New South Wales

MM Lahra, EA Limnios, TR Hogan, RL  
Kundu, J El Nasser, S Ray, M Shoushtari  
Microbiology Department, New South Wales  
Health Pathology, The Prince of Wales Hospital  
Barker Street, Randwick NSW 2031  
Telephone: +61 2 9382 9084  
Facsimile: +61 2 9382 9310  
Email: [monica.lahra@health.nsw.gov.au](mailto:monica.lahra@health.nsw.gov.au)

M Maley, R Porritt  
Department of Microbiology and Infectious  
Diseases, New South Wales Health Pathology,  
Liverpool Hospital, Locked Mail Bag 7090,  
Liverpool BC NSW 1871  
Telephone: +61 8738 5124

Facsimile: +61 2 8738 5129

Email: [Joanne.Mercer@sswahs.nsw.gov.au](mailto:Joanne.Mercer@sswahs.nsw.gov.au) or  
[Robert.Porritt@sswahs.nsw.gov.au](mailto:Robert.Porritt@sswahs.nsw.gov.au)

### Northern Territory

R Baird, K Freeman  
Microbiology Department, Territory Pathology,  
Royal Darwin Hospital, Rocklands Drive, Tiwi  
NT 0810  
Telephone: +61 8 8922 8167  
Facsimile: +61 8 8922 7788  
Email: [rob.baird@nt.gov.au](mailto:rob.baird@nt.gov.au)

### Queensland

G Robertson, H Smith, V Hicks, A. Jennison  
Public Health Microbiology, Queensland  
Health Forensic and Scientific Services, 39  
Kessels Road, Coopers Plains Qld 4108  
Telephone: +61 7 3096 2825  
Facsimile: +61 7 3096 2973; +61 7 3274 9175  
Email: [Amy.Jennison@health.qld.gov.au](mailto:Amy.Jennison@health.qld.gov.au)

### South Australia

I Bastian, A Lawrence, J Holds  
SA Pathology, Royal Adelaide Hospital Site,  
Microbiology and Infectious Diseases, Royal  
Adelaide Hospital, North Terrace, Adelaide,  
SA 5000  
Telephone: +61 8 8222 3335  
Facsimile: +61 8 2223543  
Email: [andrew.lawrence@health.sa.gov.au](mailto:andrew.lawrence@health.sa.gov.au)

### Tasmania

L Cooley, B McEwan  
Department of Microbiology and Infectious  
Diseases, Royal Hobart Hospital, 48 Liverpool  
Street, Hobart Tasmania 7000  
Telephone: +61 3 6222 8656  
Email: [belinda.mcewan@dhhs.tas.gov.au](mailto:belinda.mcewan@dhhs.tas.gov.au)

### Victoria

B Howden, K Stevens  
Microbiological Diagnostic Unit Public Health  
Laboratory, Department of Microbiology and

Immunology, The Peter Doherty Institute,  
The University of Melbourne, Parkville  
Victoria 3052  
Telephone: +61 3 8344 5713  
Facsimile: +61 3 8344 7833  
Email: [kerries@unimelb.edu.au](mailto:kerries@unimelb.edu.au)

## Western Australia

D Speers, J Bew  
Department of Microbiology, QEII Medical  
Centre, PP Block Level 5, PathWest Laboratory  
Medicine WA, Hospital Avenue, Nedlands,  
WA 6009  
Telephone: +61 8 6383 4501  
Facsimile: +61 8 9382 8046  
Email: [jane.bew@health.wa.gov.au](mailto:jane.bew@health.wa.gov.au)

## Author details

Monica M Lahra<sup>1,2</sup>  
Tiffany R Hogan<sup>1</sup>

1. Neisseria Reference Laboratory and World Health Organisation Collaborating Centre for STI and AMR, Sydney. Department of Microbiology, New South Wales Health Pathology, The Prince of Wales Hospital, Randwick, 2031, NSW Australia.
2. School of Medical Sciences, Faculty of Medicine, The University of New South Wales, NSW, 2052 Australia

## Corresponding author

Professor Monica Lahra,

Director, Microbiology Department, Neisseria Reference Laboratory and WHO Collaborating Centre for STI and AMR, New South Wales Health Pathology, Level 4, Campus Centre, The Prince of Wales Hospital, Randwick NSW, 2031.

Email: [monica.lahra@health.nsw.gov.au](mailto:monica.lahra@health.nsw.gov.au)

## References

1. National Neisseria Network. Meningococcal

isolate surveillance Australia 1994. *Commun Dis Intell.* 1995;19:286–9.

2. Australian Government Department of Health. National Notifiable Disease Surveillance System. [Internet.] Canberra: Australian Government Department of Health; 2019. Available from: <http://www9.health.gov.au/cda/source/cda-index.cfm>.
3. National Notifiable Diseases Surveillance System (NNDSS). Number of notifications of Meningococcal disease (invasive), received from state and territory health authorities in the period of 1991 to 2012 and year-to-date notifications for 2014. [Internet.] Canberra: Australian Government Department of Health; 2014.
4. Australian Government Department of Health. Invasive meningococcal disease. [Internet.] Canberra: Australian Government Department of Health; 2019. Available from: <https://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-meningococcal-W.htm>.
5. Communicable Diseases Network Australia (CDNA). Invasive meningococcal disease: CDNA national guidelines for public health units. [Internet.] Canberra: Australian Government Department of Health; 2015. Available from: <http://www.health.gov.au/internet/main/publishing.nsf/content/cdna-song-imd.htm>.
6. Abad R, López EL, Debbag R, Vázquez JA. Serogroup W meningococcal disease: global spread and current affect on the Southern Cone in Latin America. *Epidemiol Infect.* 2014;142(12):2461–70.
7. Ladhani SN, Beebeejaun K, Lucidarme J, Campbell H, Gray S, Kaczmarski E. Increase in endemic *Neisseria meningitidis* capsular group W sequence type 11 complex associated with severe invasive disease in England and Wales. *Clin Infect Dis.* 2015;60(4):578–85.

8. Chiu C, Dey A, Wang H, Menzies R, Deeks S, Mahajan D et al. Vaccine preventable diseases in Australia, 2005 to 2007. *Commun Dis Intell Q Rep*. 2010;34(Suppl):S1–167.
9. Australian Government Department of Health. Australian Immunisation Handbook: Meningococcal disease. [Internet.] Canberra: Australian Government Department of Health; 2015. Available from: <http://www.health.gov.au/internet/immunise/publishing.nsf/Content/immunise-meningococcal>
10. Araya P, Fernández J, Del Canto F, Seoane M, Ibarz-Pavón AB, Barra G et al. *Neisseria meningitidis* ST-11 clonal complex Chile, 2012. *Emerg Infect Dis*. 2015;21(2):339–41.
11. Bröker M, Jacobsson S, Kuusi M, Pace D, Simões MJ, Skoczynska A et al. Meningococcal serogroup Y emergence in Europe: update 2011. *Hum Vaccin Immunother*. 2012;8(12):1907–11.