

Communicable Diseases Surveillance

Mycoplasma pneumoniae

Atypical pneumonia due to *Mycoplasma pneumoniae* is an acute, febrile illness of the lower respiratory tract. Transmission of the organism is by the inhalation of droplets produced by coughing, or by direct contact with an infected person. The incubation period is between one and four weeks, and the infection may be asymptomatic, particularly in children under five years of age. Clinical manifestations vary from a mild afebrile pharyngitis to atypical pneumonia in up to 30% of cases. Onset is insidious with signs and symptoms including headache, malaise, cough, sore throat and occasionally pleuritic chest pain. Antimicrobial therapy is not required for an upper respiratory tract infection. Whilst pneumonia is usually self-limiting, appropriate treatment with erythromycin or tetracycline can shorten the the course of the illness, which may last from several days to a month or more. However *Mycoplasma pneumoniae* can be cultured from the sputum of infected individuals for weeks to months following effective treatment, and may therefore serve as a source of infection for others. The organism has a worldwide distribution, and whilst most cases appear to be sporadic, epidemics do occur, particularly in closed environments such as in the family setting and in institutions.

The number of reports received by the CDI Virology and Serology Laboratory Reporting Scheme has continued to rise since early 1996 (Figure 1). For 1996 a total of 1,010 reports was received. More females were reported than males, with a male:female ratio of 1:1.3 (Figure 2). The predominance of females was particularly marked in the 25 - 44 years age group.

National Notifiable Diseases Surveillance System

The NNDSS is conducted under the auspices of the Communicable Diseases Network Australia New Zealand. The system coordinates the national surveillance of more than 40 communicable diseases or disease groups endorsed by the National Health and Medical Research Council (NHMRC). Notifications of these diseases are made to State and Territory health authorities under the provisions of their respective public health legislations. De-identified core unit data are supplied fortnightly for collation, analysis and dissemination. For further information, see CDI 1997;21:5.

Correction: Tables 1 and 2 in the previous issue, CDI 1997;21:224 were incorrect, and represent data for the period 11 to 24 June 1997.

Reporting period 23 July to 5 August 1997

There were 1,931 notifications received for this two week period (Tables 1, 2 and 3). The numbers of reports for selected diseases have been compared with historical data for corresponding periods in the previous three years (Figure 4).

A total of 3,618 notifications of pertussis with onset in 1997 has been received so far. This is higher than any corresponding period since the establishment of the scheme

Figure 1. *Mycoplasma pneumoniae* laboratory reports, 1993 to 1997, by month of specimen collection




Figure 2. *Mycoplasma pneumoniae* laboratory reports, 1996, by age group and sex




Figure 3. *Haemophilus influenzae* type b infection notifications, 1991 to 1997, by month of onset




Table 1. Notifications of diseases preventable by vaccines recommended by the NHMRC for routine childhood immunisation, received by State and Territory health authorities in the period 23 July to 5 August 1997

Disease ^{1,2}	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	This period 1997	This period 1996	Year to date 1997	Year to date 1996
Diphtheria	0	0	0	0	0	0	0	0	0	0	0	0
<i>Haemophilus influenzae</i> type b	0	0	0	0	0	0	1	1	2	1	32	37
Measles	3	4	0	3	0	1	5	13	29	14	328	272
Mumps	0	1	1	NN	0	0	0	3	5	8	115	68
Pertussis	3	46	1	90	38	2	29	22	231	94	4271	1808
Rubella	5	2	0	14	4	0	12	3	40	54	763	1565
Tetanus	0	1	0	0	0	0	0	0	1	0	7	1

NN. Not Notifiable

1. No notifications of poliomyelitis have been reported since 1986.

2. Totals comprise data from all States and Territories. Cumulative figures are subject to retrospective revision, so there may be discrepancies between the number of new notifications and the increment in the cumulative figure from the previous period.

Table 2. Notifications of other diseases received by State and Territory health authorities in the period 23 July to 5 August 1997

Disease ^{1,2}	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	This period 1997	This period 1996	Year to date 1997	Year to date 1996
Arbovirus Infection (NEC) ³	0	0	1	0	0	0	0	0	1	1	108	40
Barmah Forest virus infection	0	4	-	4	0	0	0	-	8	18	494	668
Campylobacteriosis ⁴	3	-	6	227	73	8	107	36	460	451	6818	7065
Chlamydial infection (NEC) ⁵	12	NN	12	122	0	24	49	42	261	362	4840	5028
Dengue	0	0	0	0	0	-	0	1	1	2	193	26
Donovanosis	0	NN	6	0	NN	0	0	0	6	2	23	32
Gonococcal infection ⁶	1	19	19	33	0	0	16	28	116	174	2713	2490
Hepatitis A	2	13	3	46	1	0	7	3	75	68	2022	1466
Hepatitis B incident	0	0	0	0	0	0	4	0	4	6	134	143
Hepatitis C incident	0	0	0	-	0	0	-	-	0	3	8	32
Hepatitis C unspecified	7	NN	10	128	NN	11	129	6	291	307	5607	5929
Hepatitis (NEC)	0	0	0	0	0	0	0	NN	0	1	12	12
Legionellosis	0	0	0	0	1	0	0	0	1	7	100	117
Leptospirosis	0	1	0	1	0	0	0	0	2	16	78	155
Listeriosis	0	1	0	0	0	0	1	1	3	6	52	35
Malaria	3	0	0	0	2	1	4	1	11	48	493	529
Meningococcal infection	0	2	0	3	1	0	2	6	14	36	235	213
Ornithosis	0	NN	0	0	0	0	2	0	2	1	37	55
Q Fever	0	3	0	6	0	0	0	1	10	20	355	321
Ross River virus infection	0	6	2	25	1	1	2	4	41	40	6203	7352
Salmonellosis (NEC)	3	16	9	45	13	3	28	13	130	156	4685	3836
Shigellosis ⁴	0	-	2	9	4	0	3	4	22	42	533	429
Syphilis	0	8	3	13	0	2	0	1	27	85	726	944
Tuberculosis	0	1	0	8	2	0	17	1	29	29	571	648
Typhoid ⁷	0	1	0	0	0	0	2	0	3	3	46	59
Yersiniosis (NEC) ⁴	1	-	0	10	1	0	0	0	12	4	173	156

1. For HIV and AIDS, see *CDI* 1997;21:226-227. For rarely notified diseases, see Table 3.

2. Totals comprise data from all States and Territories. Cumulative figures are subject to retrospective revision so there may be discrepancies between the number of new notifications and the increment in the cumulative figure from the previous period.

3. NT: includes Barmah Forest virus.

4. NSW: only as 'foodborne disease' or 'gastroenteritis in an institution'.

5. WA: genital only.

6. NT, Qld, SA and Vic: includes gonococcal neonatal ophthalmia.


7. NSW, Vic: includes paratyphoid.

NN Not Notifiable.

NEC Not Elsewhere Classified

- Elsewhere Classified.

Figure 4. Selected National Notifiable Diseases Surveillance System reports, and historical data¹



1. The historical data are the averages of the number of notifications in 9 previous 2-week reporting periods, the corresponding periods of the last 3 years and the periods immediately preceding and following those.

Figure 5. Syphilis notifications, 1991 to 1997, by month of onset




Figure 6. Syphilis notifications, 1997, by age group and sex




Table 3. Notifications of rare¹ diseases received by State and Territory health authorities in the period 23 July to 5 August 1997

Disease ²	Total this period	Reporting States or Territories	Total notifications 1997
Brucellosis	2	Qld	20
Chancroid			1
Cholera			2
Hydatid infection	3	NSW, Vic	24
Leprosy			7

1. Fewer than 60 cases of each of these diseases were notified each year during the period 1988 to 1996.
2. No notifications have been received during 1997 for the following rare diseases: botulism, lymphogranuloma venereum, plague, rabies, yellow fever, or other viral haemorrhagic fevers.

A total of 3,618 notifications of pertussis with onset in 1997 has been received so far. This is higher than any corresponding period since the establishment of the scheme in 1990. There were 231 notifications of pertussis this period. Notifications are expected to increase over the spring and summer months, and the total notifications for 1997 is likely to be the highest recorded by this scheme.

There have been 32 notifications of *Haemophilus influenzae* type b infection (Hib) for the year to date. The number of notifications has declined following the introduction of conjugate Hib vaccines in 1992 (Figure 3).

A decrease in the number of notifications of syphilis has been observed since 1995 (Figure 5). For 1997 the male:female ratio was 1.2:1. However, there was a predominance of females in the 15 - 19 years age group (Figure 6).

National Influenza Surveillance, 1997

Three types of data are included in National Influenza Surveillance, 1997. These are sentinel general practitioner surveillance conducted by the Australian Sentinel Practice Research Network, Department of Human Services, Victoria, Department of Health, New South Wales and Department of Health and Community Services, Northern Territory; laboratory surveillance data from the Communicable Diseases Intelligence Virology and Serology Laboratory Reporting Scheme, LabVISE, and the World Health Organization Collaborating Centre for Influenza Reference and Research; and absenteeism surveillance conducted by Australia Post. For further information about these schemes, see *CDI 1997; 21:126*.

Overall influenza activity continued to rise this fortnight, although the sentinel general practitioner consultation rate recorded in the Northern Territory declined. Sixty per cent of reports this period were for influenza B. Reports of influenza A have however, increased.

Laboratory Surveillance

A total of 323 reports of influenza virus was recorded by the LabVISE scheme this fortnight (Figure 7). Of these 130 were for influenza A, 177 for influenza B and 16 were untyped. The epidemic of influenza B this season is

Figure 7. Laboratory reports of influenza, 1997, by type and week of specimen collection




Table 4. Australian Sentinel Practice Research Network reports, weeks 30 and 31, 1997

Condition	Week 30, to 27 July 1997		Week 31, to 3 August 1997	
	Reports	Rate per 1,000 encounters	Reports	Rate per 1,000 encounters
Chickenpox	5	0.8	17	2.3
Gastroenteritis	51	8.3	58	7.9
HIV testing (doctor initiated)	7	1.1	9	1.2
HIV testing (patient initiated)	9	1.5	11	1.5
Influenza	184	29.9	244	33.1
Measles	1	0.2	2	0.3
Pertussis	1	0.2	1	0.1
Ross River virus infection	1	0.2	6	0.8
Rubella	2	0.3	1	0.1

Table 5. Sentinel Chicken Surveillance Programme seroconversions, Western Australia, June and July 1997

	June				July			
	MVE	Kunjin	MVE & Kunjin	Flavivirus	MVE	Kunjin	MVE & Kunjin	Flavivirus
Kimberley								
Kalumburu	2		1		2			
Kununurra	1							
Fitzroy Crossing	1							
Derby	1		1					
Lombadina	1							
Broome			1		2	1	1	
Pilbara								
Karratha	1							

Sentinel Chicken Surveillance Programme

Sentinel chicken flocks are used to monitor flavivirus activity in Australia. The main viruses of concern are Murray Valley encephalitis (MVE) and Kunjin which cause the potentially fatal disease Australian encephalitis in humans. Currently 24 flocks are maintained in the north of Western Australia, ten in the Northern Territory, ten in New South Wales and ten in Victoria. The flocks in Western Australia and the Northern Territory are tested year round but those in New South Wales and Victoria are tested only from November to March, during the main risk season. Results are coordinated by the Arbovirus Laboratory in Perth and reported bimonthly. For more information see *CDI 1997;21:6*

AK Broom¹, JS Mackenzie², L Melville³, DW Smith⁴ and PI Whelan⁵

1. Department of Microbiology, The University of Western Australia
2. Department of Microbiology, The University of Queensland
3. Berrimah Agricultural Research Centre, Darwin
4. PathCentre, Perth
5. Department of Health and Community Services, Darwin

Sentinel chicken serology was carried out for all of the 24 flocks in Western Australia in June and July 1997. There

were 11 seroconversions to flaviviruses in the Kimberley and Pilbara regions in June, and 6 from the Kimberley region in July (Table 5). Five flocks of sentinel chickens from the Northern Territory were tested in June and July 1997, and during this period there were no seroconversions to flaviviruses.


LabVISE

The Virology and Serology Laboratory Reporting Scheme, LabVISE, is a sentinel reporting scheme. Twenty-one laboratories contribute data on the laboratory identification of viruses and other organisms. Data are collated and published in *Communicable Diseases Intelligence* each fortnight. These data should be interpreted with caution as the number and type of reports received is subject to a number of biases. For further information, see *CDI 1997;21:8-9*.

There were 2,131 reports received in the *CDI* Virology and Serology Laboratory Reporting Scheme this period (Tables 6 and 7).

The number of Ross River virus reports has declined after peaking in March. There were 51 laboratory reports of Ross River virus this fortnight with 50% of reports received from Queensland and 43% from Western Australia.

Figure 9. Rhinovirus laboratory reports, 1991 to 1996 average and 1997, by month of specimen collection



Forty-six reports of rhinovirus were received this period. Ninety-one per cent of reports were for the 1 - 4 years age group. The number of reports received this year is low compared to previous years (Figure 9).

Six hundred and ninety-seven reports of respiratory syncytial virus were received this period. Ninety-three per cent of reports were for children below the age of five years. The number of reports received this year is consistent with that of previous years (Figure 10).

One hundred and four reports of rotavirus were received this period for 54 males and 48 females (2 sex not stated). Eighty-six per cent of reports were for children under five years of age. The number of reports was lower than average for the month of June (Figure 11).

Figure 10. Respiratory syncytial virus laboratory reports, 1994 to 1996 average and 1997, by month of specimen collection




Figure 11. Rotavirus laboratory reports, 1992 to 1996 average and 1997, by month of specimen collection




Table 6. Virology and serology laboratory reports by State or Territory¹ for the reporting period 17 to 30 July 1997, historical data², and total reports for the year

	State or Territory ¹								Total this fortnight	Historical data ²	Total reported in CDI in 1997
	ACT	NSW	NT	Qld	SA	Tas	Vic	WA			
Measles, mumps, rubella											
Measles virus				1				2	3	2.3	40
Mumps virus				1				4	5	1.5	28
Rubella virus	3			1			3	1	8	14.8	412
Hepatitis viruses											
Hepatitis A virus	7	1		15	2		2	15	42	11.5	541
Hepatitis D virus					1				1	0.7	15
Arboviruses											
Ross River virus			1	22	1		2	25	51	18.5	2,005
Barmah Forest virus				5				1	6	6.7	193
Dengue not typed				2				10	12	1.2	54
Kunjin virus								1	1	0	7
Adenoviruses											
Adenovirus type 1					1		1		2	1.3	19
Adenovirus type 41								2	2	0	3
Adenovirus not typed/pending	6	2		29	10		2	6	55	47.7	628

Table 6. Virology and serology laboratory reports by State or Territory¹ for the reporting period 17 to 30 July 1997, historical data², and total reports for the year, continued

	State or Territory ¹								Total this fortnight	Historical data ²	Total reported in <i>CDI</i> in 1997
	ACT	NSW	NT	Qld	SA	Tas	Vic	WA			
Herpes viruses											
Herpes virus type 6								1	1	0	4
Cytomegalovirus	5	8		26	7		10	18	74	62.8	788
Varicella-zoster virus	5	2	1	18	9		13	28	76	49.5	936
Epstein-Barr virus	7	12	1	14	30	1	5	24	94	81.3	1,748
Other DNA viruses											
Molluscum contagiosum								1	1	0.2	7
Contagious pustular dermatitis								2	2	0	2
Parvovirus					4	1	10	1	16	8.2	252
Picornavirus family											
Coxsackievirus A9	3								3	0	6
Coxsackievirus A16	2								2	0	10
Echovirus type 4					1				1	0	1
Echovirus type 5	1								1	0	6
Echovirus type 6					1				1	0	1
Echovirus type 9	1								1	0.7	2
Echovirus not typed/pending					1				1	0	3
Rhinovirus (all types)		9		16	2		1	18	46	30.3	414
Enterovirus not typed/pending				24				34	58	29.8	430
Ortho/paramyxoviruses											
Influenza A virus	1	31		25	9		40	19	125	181.8	477
Influenza A virus H3N2				1		1	3		5	7.3	7
Influenza B virus	2	14	1	48	3	3	44	62	177	22.2	530
Influenza virus - typing pending					16				16	0.3	230
Parainfluenza virus type 1			1						1	14.3	42
Parainfluenza virus type 2				7	5			1	13	7.5	93
Parainfluenza virus type 3	1	3	1	14	6		4	8	37	34.7	503
Parainfluenza virus typing pending					4				4	2.2	189
Respiratory syncytial virus	54	189		206	57	15	153	23	697	525.8	2,791
Paramyxovirus (unspecified)							1		1	0.8	13
Other RNA viruses											
HTLV-1			1						1	0.5	10
Rotavirus	6	5		24	2	35	32		104	127.3	712
Norwalk agent						2			2	1	67
<i>Chlamydia trachomatis</i> not typed	52	10		30	16	3	4	129	244	139.7	3,275
<i>Chlamydia psittaci</i>							1		1	3	47
<i>Chlamydia</i> species				1					1	0.7	22
<i>Mycoplasma pneumoniae</i>	4	10	2	23	7	2	8	12	68	27.3	1,140
<i>Coxiella burnetii</i> (Q fever)		5		8				5	18	8.2	232
<i>Rickettsia australis</i>				1					1	1.2	13
<i>Rickettsia tsutsugamushi</i>				3					3	1.3	20
<i>Bordetella pertussis</i>	2	1		10			9	19	41	11	1,134
<i>Legionella pneumophila</i>								1	1	0.7	16
<i>Cryptococcus</i> species	1								1	0.3	14
<i>Leptospira pomona</i>				1					1	0	12
<i>Leptospira</i> species				1					1	2.2	6
<i>Toxoplasma gondi</i>							1		1	2.5	1
TOTAL	163	302	9	553	217	28	354	505	2,131	1,492.80	20,151

1. State or Territory of postcode, if reported, otherwise State or Territory of reporting laboratory.

2. The historical data are the averages of the numbers of reports in 6 previous 2 week reporting periods, the corresponding periods of the last 2 years and the periods immediately preceding and following those.

Table 7. Virology and serology laboratory reports by contributing laboratories for the reporting period 17 to 30 July 1997

State and Territory	Laboratory	Reports
Australian Capital Territory	The Canberra Hospital, Canberra	187
New South Wales	Institute of Clinical Pathology & Medical Research, Westmead	31
	New Children's Hospital, Westmead	226
	Royal Prince Alfred Hospital, Camperdown	18
Queensland	Queensland Medical Laboratory,	172
	West End State Health Laboratory, Brisbane	389
South Australia	Institute of Medical and Veterinary Science, Adelaide	214
Tasmania	Northern Tasmanian Pathology Service, Launceston	23
	Royal Hobart Hospital, Hobart	1
Victoria	Commonwealth Serum Laboratories, Melbourne	9
	Microbiological Diagnostic Unit, University of Melbourne	4
	Monash Medical Centre, Melbourne	51
	Royal Children's Hospital, Melbourne	173
	Victorian Infectious Diseases Reference Laboratory, Fairfield	121
Western Australia	PathCentre, Virology, Perth	512
TOTAL		2131