



Australian Government
Department of Health
and Aged Care



Australian
Centre for
Disease
Control

2024 • Volume 48

Communicable Diseases Intelligence

Australian Group on Antimicrobial Resistance surveillance outcome programs – bloodstream infections and antimicrobial resistance patterns from patients less than 18 years of age, January 2020 – December 2021

Anita Williams, Geoffrey W Coombs, Jan Bell, Denise A Daley, Shakeel Mowlaboccus,
Penelope A Bryant, Anita J Campbell, Louise Cooley, Jon Iredell, Adam D Irwin, Alison Kesson,
Brendan McMullan, Morgyn S Warner, Phoebe Williams, Christopher C Blyth

Communicable Diseases Intelligence

Communicable Diseases Intelligence (CDI) is a peer-reviewed scientific journal published by the Health Security & Emergency Management Division, Department of Health and Aged Care.

The journal aims to disseminate information on the epidemiology, surveillance, prevention and control of communicable diseases of relevance to Australia.

© 2024 Commonwealth of Australia as represented by the Department of Health and Aged Care

ISSN: 2209-6051 Online

This journal is indexed by Index Medicus and Medline.

Creative Commons Licence – Attribution-NonCommercial-NoDerivatives CC BY-NC-ND



This publication is licensed under a Creative Commons Attribution-Non-Commercial NoDerivatives 4.0 International Licence from <https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode> (Licence). You must read and understand the Licence before using any material from this publication.

<https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode> (Licence). You must read and understand the Licence before using any material from this publication.

Restrictions

The Licence does not cover, and there is no permission given for, use of any of the following material found in this publication (if any):

- the Commonwealth Coat of Arms (by way of information, the terms under which the Coat of Arms may be used can be found at www.pmc.gov.au/resources/commonwealth-coat-arms-information-and-guidelines);
- any logos (including the Department of Health and Aged Care's logo) and trademarks;
- any photographs and images;
- any signatures; and
- any material belonging to third parties.

Disclaimer

Opinions expressed in *Communicable Diseases Intelligence* are those of the authors and not necessarily those of the Australian Government Department of Health and Aged Care or the Communicable Diseases Network Australia. Data may be subject to revision.

Enquiries

Enquiries regarding any other use of this publication should be addressed to the CDI Editor at: cdi.editor@health.gov.au

Communicable Diseases Network Australia

Communicable Diseases Intelligence contributes to the work of the Communicable Diseases Network Australia. www.health.gov.au/cdna

Editor

Christina Bareja

Deputy Editor

Simon Petrie

Design and Production

Lisa Thompson

Editorial Advisory Board

David Durrheim, Mark Ferson, Clare Huppertz, John Kaldor, Martyn Kirk, Meru Sheel and Stephanie Williams

Contacts

CDI is produced by:

Health Security & Emergency Management Division
Australian Government Department of Health and Aged Care

GPO Box 9848, (MDP 6)
CANBERRA ACT 2601

www.health.gov.au/cdi

cdi.editor@health.gov.au

Submit an Article

You are invited to submit your next communicable disease related article to *Communicable Diseases Intelligence* (CDI) for consideration. More information regarding CDI can be found at: www.health.gov.au/cdi.

Further enquiries should be directed to: cdi.editor@health.gov.au.

Australian Group on Antimicrobial Resistance surveillance outcome programs – bloodstream infections and antimicrobial resistance patterns from patients less than 18 years of age, January 2020 – December 2021

Anita Williams, Geoffrey W Coombs, Jan Bell, Denise A Daley, Shakeel Mowlaboccus, Penelope A Bryant, Anita J Campbell, Louise Cooley, Jon Iredell, Adam D Irwin, Alison Kesson, Brendan McMullan, Morgyn S Warner, Phoebe Williams, Christopher C Blyth

Abstract

From 1 January 2020 to 31 December 2021, thirty-eight institutions across Australia submitted data to the Australian Group on Antimicrobial Resistance (AGAR) from patients aged < 18 years (AGAR-Kids). Over the two years, 1,679 isolates were reported from 1,611 patients. This AGAR-Kids report aims to describe the population of children and adolescents with bacteraemia reported to AGAR and the proportion of resistant isolates.

Overall, there were 902 gram-negative isolates reported: 800 Enterobacterales, 61 *Pseudomonas aeruginosa* and 41 *Acinetobacter* spp. Among the Enterobacterales, 12.9% were resistant to third generation cephalosporins; 11.6% to gentamicin/tobramycin; and 11.2% to piperacillin-tazobactam. In total, 14.5% of Enterobacterales were multi-drug resistant (MDR). Only 3.3% of *P. aeruginosa* were resistant to carbapenems and 4.9% were MDR. Resistance in *Acinetobacter* spp was uncommon.

Of 607 *Staphylococcus aureus* isolates, 12.9% were methicillin-resistant (MRSA). Almost half of *S. aureus* isolates from the Northern Territory were MRSA. In *S. aureus*, resistance to erythromycin was 13.2%; 12.4% to clindamycin; and 5.3% to ciprofloxacin. Resistance to all antibiotics tested was higher in MRSA. Overall, 6.5% of *S. aureus* were MDR, of which 65% were MRSA.

Almost three-quarters of the 170 *Enterococcus* spp. reported were *E. faecalis*, and half were from patients < 1 year old. Ampicillin resistance in enterococci was 19.6%. Eight isolates were vancomycin resistant and three isolates were teicoplanin resistant. Five *E. faecium* isolates were classified as MDR.

This AGAR-Kids report highlights clear differences in the geographic distribution of pathogens and resistance profiles across Australia.

Keywords: Australian Group on Antimicrobial Resistance (AGAR); antimicrobial resistance surveillance; paediatrics; bacteraemia; Enterobacterales; *Staphylococcus aureus*; *Enterococcus*

Introduction

The Australian Group on Antimicrobial Resistance (AGAR) produces annual reports from the whole of the Australian population. However, previous analysis of the AGAR data, comparing adult (> 18 years) and paediatric (\leq 18 years) bacteraemia, suggests there are lower rates of antimicrobial-resistant organisms isolated in children, with different phenotypes and lower mortality rates.

In 2018, the World Society for Paediatric Infectious Diseases (WSPID) declared that antimicrobial resistance (AMR) surveillance programs should present neonatal- and paediatric-specific data to assist with strengthening knowledge.¹ Currently, age-specific data are not routinely reported in the majority of AMR surveillance programs, making paediatric-specific interventions difficult.² By monitoring paediatric bacteraemia, we are able to provide insight not only into the dynamic aetiology of bloodstream infections, but also into the impact of vaccine programs, and to inform strategies aimed at targeting invasive infections.³

The key influences on bacteraemia incidence are age, vaccination coverage and exposure to invasive procedures.³ In previous studies, neonates were often over-represented, reflecting the immaturity of the immune system and the use of invasive devices (e.g., intravenous catheters). Various reports from Australia and Europe suggest there are differences in the AMR burden of various organisms, not only between adults and children, but within different age groups among children. For example, gram-negative multidrug resistant (MDR; resistant to three or more antimicrobial classes) organisms were previously found to disproportionately impact children, as demonstrated by a higher odds of death in children with bacteraemia secondary to an extended spectrum beta-lactamase (ESBL) containing organism vs non-ESBL bacteraemia when compared to the same ratio in adults.⁴ In a European study, isolates from children < 1 year of age had less AMR than those isolated from children \geq 1 year old.² In a Scottish study, instances of bacteraemia in children < 1 year old were more likely to be healthcare associated, whilst bacteraemia in children aged 1–15 years were more often community associated.⁵ These findings impact the empiric treatment guidelines and stewardship initiatives, thereby highlighting the value of paediatric-specific reporting of bacteraemia.

Methods

Participants

Thirty-eight laboratories participating in AGAR who reported data from patients < 18 years old (0–17 years of age inclusive); 35 laboratories in each year respectively. AGAR collects data from 24 of the 29 principal referral centres and from all tertiary paediatric centres (8 hospitals) in Australia, as well as from seven public acute group A hospitals, three private group A hospitals, two private group B hospitals and all public group C hospitals in the north-west of Western Australia.⁶

Collection period

From 1 January 2020 to 31 December 2021, participating laboratories submitted all isolates from unique patient episodes of bacteraemia for *S. aureus* and *Enterococcus* spp., and up to 200 isolates per year for Enterobacterales, *P. aeruginosa*, and *Acinetobacter* spp.

An episode was defined as a clinical event associated with a positive blood culture, irrespective of the number of bacterial species identified. A new episode of bacteraemia in the same patient was recorded if the blood culture was collected more than two weeks after the initial positive culture. An episode was defined as community-onset if the first positive blood culture of the episode was collected 48 hours or less after hospital admission, and as hospital-onset if collected more than 48 hours after admission.

Laboratory methods

Isolates were identified to the species level by the participating laboratories using matrix-assisted laser desorption ionization (MALDI) [MALDI Biotyper (Bruker Daltonics, United States of America [USA]) or Vitek-MS (bioMérieux, France) or using Vitek2[®] (bioMérieux). Antimicrobial susceptibility testing was performed using the Vitek2[®] (bioMérieux) or BD Phoenix[™] (Becton Dickinson, USA) automated microbiology systems according to the manufacturer's instructions.

Data collection and analysis

Data were collected on age, sex, dates of admission and discharge, and mortality at seven and 30 days from date of blood culture collection. To avoid interpretive bias, no attempt was made to assign attributable mortality.

The AMR for R package (v2.0) was used to transform minimum inhibitory concentration data as per EUCAST 2022 (v12) breakpoints.^{7,8} Multi-drug resistance (MDR) was defined as resistance to one or more agents in three or more antimicrobial categories.⁹ Descriptive statistics for the population and isolates for the overall population and per year were stratified by age, sex, and state/territory where appropriate. Categorical data was assessed using the chi-square or Fisher's exact test. Continuous data was assessed using the Student t-test or Mann-Whitney U test. Proportions were not calculated where fewer than ten isolates in a category were tested.

Ethics

Approval to conduct the prospective data collection was given by the research ethics committee associated with each participating healthcare facility.

Results

Patient characteristics

Overall, there were 1,679 isolates from 1,611 bacteraemic episodes in 1,611 patients: 826 episodes with 856 isolates in 2020, and 785 episodes with 823 isolates in 2021. The most frequently reported species were *S. aureus* (n: 607/1,679; 36.2%) and *Escherichia coli* (n: 378/1,679; 22.5%). In all states and mainland territories, *S. aureus* was the most frequently reported species, followed by *E. coli*, except in Queensland where non-typhoidal *Salmonella* was more frequent. For patients aged < 1 year, the most common species was *E. coli*, whereas in patients aged ≥ 1 year, the most common species was *S. aureus* (Table 1).

There was an over-representation of episodes in patients aged < 1 year (n: 623/1,611; 38.7%), and 15.1% of all episodes were in neonates (n: 244). More bacteraemia episodes affected males (n: 976/1,611; 60.6%). The median age of patients with *S. aureus* bacteraemia was older than for enterococcal or gram-negative bacteraemia.

Most bacteraemic episodes were community-onset (69.0%); however, enterococcal episodes were more often hospital-onset. The proportion of hospital-onset episodes was highest in the neonatal age group and decreased with age; the median age for community-onset bacteraemia was 3 years (interquartile range, IQR: 0–10 years), whereas the median age for hospital-onset bacteraemia was < 12 months (IQR: 0–5 years). Tasmania had the highest proportion of community-onset episodes (85% of all episodes) with Victoria having the lowest proportion (58% of episodes).

Overall, the mean length of stay following blood culture collection was 17.0 days (standard deviation, SD: 21.5) and the median was 10 days (IQR: 6–21 days). At 30 days, 21.7% of all children with bacteraemia (n = 350/1,611) were still admitted to hospital.

Over the two-year period, bone and joint infections were the most frequently reported clinical manifestations (n: 255/1,679; 15.2%), followed by device-related infections without metastatic focus (n: 210/1,679; 12.5%). Community-onset bacteraemic episodes were more frequently associated with an osteoarticular focus, whereas hospital-onset episodes were more frequently device-related or in patients with febrile neutropenia.

Table 1: Characteristics of patients aged < 18 years with a bacteraemic event reported to AGAR, per survey, January 2020 – December 2021

Category	Characteristic	Enterococcus spp. (N = 167)		S. aureus (N = 606)		Gram-negative bacteria (N = 867)		Total patients (N = 1,611) ^a	
		n	%	n	%	n	%	n	%
Age (years)	Median	<1	—	6	—	1	—	2	—
	IQR ^b	0–4	—	1–11.8	—	0–7	—	0–10	—
Age group	≤ 28 days	38	22.8	51	8.4	160	18.5	244	15.1
	29–90 days	26	15.6	51	8.4	104	12.0	173	10.7
	91–364 days	24	14.4	45	7.4	139	16.0	206	12.8
	1–4 years	38	22.8	133	21.9	198	22.8	359	22.3
	5–17 years	41	24.6	326	53.8	266	30.7	629	39.0
Sex	Female	60	35.9	208	34.3	377	43.5	635	39.4
	Male	107	64.1	398	65.7	490	56.5	976	60.6
Jurisdiction ^c	ACT	3	1.8	20	3.3	22	2.5	45	2.8
	NSW	68	40.7	172	28.4	290	33.4	516	32.0
	NT	2	1.2	29	4.8	41	4.7	72	4.5
	Qld	10	6.0	88	14.5	116	13.4	213	13.2
	SA	13	7.8	57	9.4	63	7.3	130	8.1
	Tas.	4	2.4	15	2.5	20	2.3	39	2.4
	Vic.	56	33.5	152	25.1	239	27.6	438	27.2
	WA	11	6.6	73	12.0	76	8.8	158	9.8
	Median	12.5	—	11	—	10	—	10	—
	IQR ^b	7.8–34.3	—	7–18	—	5–21	—	6–21	—
30-day outcome	Died	6	3.6	6	1.0	42	4.8	53	3.3
	Survived	150	89.8	515	85.0	702	81.0	1,340	83.2
	Unknown	11	6.6	85	14.0	123	14.2	218	13.5
Onset	Community	70	41.9	477	78.7	583	67.2	1,112	69.0
	Hospital	97	58.1	129	21.3	284	32.8	499	31.0
Device-related infection	Yes	72	43.1	108	17.8	212	24.5	379	23.5
	No	90	53.9	473	78.1	600	69.2	1,147	71.2
	Unknown	5	3.0	25	4.1	55	6.3	85	5.3
Polymicrobial		60	35.9	43	7.1	100	11.5	174	10.8

^a The numbers in each organism group may not add up to the total column, as some patients have polymicrobial bacteraemic episodes and are reported in each respective column to their episode, but only counted once in the total number of patients.

^b IQR: interquartile range.

^c ACT: Australian Capital Territory; NSW: New South Wales; NT: Northern Territory; Qld: Queensland; SA: South Australia; Tas.: Tasmania; Vic.: Victoria; WA: Western Australia.

Overall, 23.5% of patients had a device-related bacteraemic episode (n: 379/1,611). The proportion was higher in 2021 (n: 213; 27.4%) than in 2020 (n: 166; 22.2%). The median length of stay was longer for patients with a device-related episode than for those patients without a device-related episode. More device-related episodes were reported with gram-negative isolates. Overall, 13% of episodes that were MDR were associated with a device-related infection.

Ten percent of patients had more than one isolate reported in a bacteraemia episode (174/1,611; 10.8%). The largest proportion of polymicrobial episodes were reported in children < 12 months (72/174; 41.4%).

At seven days after blood culture collection, 2.6% of patients had died (n: 42); the proportion of patients who died increased to 3.3% at 30 days (n: 53). The median age of patients who died was younger than for survivors (X^2 : 12.5; p : < 0.001). More than half of the deaths occurred in patients \leq 28 days of age (27/53; 51%), including ten neonates with community-onset bacteraemia.

Gram-negative isolates

Overall, there were 902 gram-negative isolates reported from 867 bacteraemia episodes in patients aged < 18 years in 2020–2021; 800 Enterobacterales (88.7%), 61 *Pseudomonas aeruginosa* (6.8%) and 41 (4.5%) *Acinetobacter* spp. The most frequently reported Enterobacterales were *E. coli* and *Klebsiella pneumoniae* complex.

The largest proportion of Enterobacterales were from New South Wales, followed by Victoria. The highest proportion of isolates were from patients < 12 months old, and from males. Enterobacterales episodes were more frequently community-onset, and 14% of episodes were polymicrobial (Table 1). Overall, 18% of the Enterobacterales were MDR.

The proportion of Enterobacterales resistant to the aminoglycosides (gentamicin/tobramycin) was 11.6% (n: 92/793; 95% CI: 9.5–14.0%). Victoria reported the highest proportion of aminoglycoside-resistant isolates, followed by South Australia and New South Wales. No aminoglycoside-resistant isolates were reported in Tasmania. Resistant isolates were more frequently reported in children aged 1–4 years of age. Isolates from patients with a hospital-onset episode were more frequently reported as resistant than those with a community-onset episode (Table 2).

The proportion of Enterobacterales resistant to piperacillin-tazobactam was 11.2% (88/784; 95% CI: 9.1–13.6%). Victoria reported the highest proportion of resistant isolates, followed by New South Wales. No piperacillin-tazobactam resistant isolates were reported in Tasmania and the Australian Capital Territory (Table 2).

Overall, 13.2% of Enterobacterales (101/766; 95% CI: 10.9–15.8) were reported as ciprofloxacin resistant. Isolates from patients with a hospital-onset episode were more frequently reported as ciprofloxacin resistant than those with a community-onset bacteraemia. Isolates from children aged over 1 year were more frequently resistant. Victoria reported the highest proportion of resistant isolates (Table 2).

Of the 792 Enterobacterales that had meropenem susceptibility testing performed, only two *Enterobacter cloacae* complex isolates were resistant (0.3%; 95% CI: 0.0–0.9%). The isolates reported were from a community-onset bacteraemic episode in Queensland and a hospital-onset bacteraemic episode in Victoria, both in patients aged 5–17 years.

Fourteen percent of Enterobacterales isolates were identified as MDR. MDR Enterobacterales isolates were more likely to be hospital-onset (p : < 0.01) and associated with a device-related infection (p : < 0.01). MDR Enterobacterales isolates were more frequent in patients with febrile neutropenia. Of the 41 patients who died with an Enterobacterales episode, 19.5% of the isolates were MDR (n: 8). The highest proportion of MDR Enterobacterales isolates were from Victoria (n: 59, 39.6%) and New South Wales (n: 53, 35.6%). Victoria had the highest proportion of Enterobacterales isolates that were MDR (24.8%), followed by New South Wales (20.1%) [Table 3].

There were 61 *Pseudomonas aeruginosa* reported to AGAR: 31 isolates in 2020 and 30 isolates in 2021. No isolates were reported from the Australian Capital Territory, and only one isolate was reported from Tasmania. Overall, 4.9% of *P. aeruginosa* isolates were MDR, and two isolates were carbapenem resistant (3.3%), classified as a World Health Organization (WHO) priority pathogen.

No *P. aeruginosa* isolates were reported tobramycin resistant. Overall, 19.7% of *P. aeruginosa* were piperacillin-tazobactam resistant (n: 12; 95% CI: 10.6–31.8%). All ciprofloxacin-resistant *P. aeruginosa* isolates were reported from patients aged 5–17 years. Only two *P. aeruginosa* were reported meropenem resistant (3.3%; 95% CI: 0.4–11.3%): one isolate from Queensland and one isolate from Victoria.

Table 2: Number of Enterobacterales isolates tested and proportion of isolates resistant to gentamicin/tobramycin, ceftriaxone/ceftazidime, piperacillin-tazobactam, and ciprofloxacin from children aged < 18 years as reported to AGAR, January 2020 – December 2021

Category	Characteristic	Gentamicin/tobramycin ^a			Ceftriaxone/ceftazidime ^a			Piperacillin-tazobactam ^a			Ciprofloxacin ^a						
		N	n	R%	95% CI	N	n	R%	95% CI	N	n	R%	95% CI				
Overall		793	92	11.6	9.5–14.0	793	102	12.9	10.6–15.4	784	88	11.2	9.1–13.6	766	101	13.2	10.9–15.8
Jurisdiction ^b	ACT	23	1	4.3	0.1–21.9	23	1	4.3	0.1–21.9	23	0	0	—	23	1	4.3	1.1–28.0
	NSW	264	31	11.9	8.2–16.5	264	37	14.2	10.2–19.1	260	34	13.1	9.2–17.8	250	37	14.2	7.2–15.3
	NT	37	3	8.3	1.8–22.5	37	2	5.6	0.7–18.7	36	3	8.3	1.8–22.5	35	2	5.6	—
	Qld	99	6	6.1	2.3–12.7	99	6	6.1	2.3–12.7	97	8	8.2	3.6–15.6	85	6	6.1	5.8–20.6
	SA	56	8	14.5	6.5–26.7	56	4	7.3	2.0–17.6	55	1	1.8	0.0–9.7	55	4	7.3	6.5–26.7
	Tas.	17	0	0	—	17	1	5.9	0.1–28.7	17	0	0	—	17	1	5.9	1.5–36.4
	Vic.	238	40	16.8	12.3–22.2	238	46	19.3	14.5–24.9	235	36	15.3	11.0–20.6	236	46	19.3	15.4–26.0
WA	66	3	4.6	1.0–12.9	66	5	7.7	2.5–17.0	61	6	9.8	3.7–20.2	65	5	7.7	1.7–15.0	
Age group	≤ 28 days	157	21	13.5	8.5–19.8	157	13	8.3	4.5–13.8	155	2	1.3	0.2–4.6	156	13	8.3	6.0–16.1
	29–90 days	102	10	9.9	4.9–17.5	102	11	10.9	5.6–18.7	97	12	12	6.4–20.0	98	11	10.9	4.9–17.5
	91–364 days	132	9	6.9	3.2–12.6	132	12	9.2	4.8–15.5	132	9	7	3.2–12.8	126	12	9.2	4.0–14.4
	1–4 years	173	30	17.4	12.1–24.0	173	35	20.3	14.6–27.1	170	27	15.9	10.7–22.3	160	35	20.3	13.6–26.4
	5–17 years	236	22	9.4	6.0–13.9	236	31	13.3	9.2–18.4	230	38	16.5	12.0–22.0	226	31	13.3	10.6–20.4
Onset	Community	540	39	7.2	5.2–9.7	540	44	8.1	6.0–10.8	534	38	7.1	5.1–9.6	513	44	8.1	7.7–13.1
	Hospital	253	53	20.9	16.1–26.5	253	58	22.9	17.9–28.6	250	50	20	15.2–25.5	253	58	22.9	14.7–24.8

^a N: number of isolates tested; n: number of resistant isolates; R%: proportion of resistant isolates; 95% CI: 95% confidence interval.

^b ACT: Australian Capital Territory; NSW: New South Wales; NT: Northern Territory; Qld: Queensland; SA: South Australia; Tas.: Tasmania; Vic.: Victoria; WA: Western Australia.

Table 3: Characteristics of patients aged < 18 years with a bacteraemic event reported to AGAR for which one or more isolate was multidrug resistant (MDR), per survey, January 2020 – December 2021

Category	Characteristic	Enterococcus spp. (n = 4/163) ^a		S. aureus (n = 34/572) ^a		Gram-negative bacteria (n = 118/749) ^a		Total patients (n = 156/1,484) ^{a,b}	
		n	%	n	%	n	%	n	%
Age (years)	Median	<1	—	<1	—	2	—	1	—
	IQR ^c	0–0.3	—	0–3.8	—	0–8	—	0–8	—
Age group	≤ 28 days	1	2.6	5	9.8	18	11.3	24	9.6
	29–90 days	1	3.8	7	13.7	12	11.5	20	11.0
	91–364 days	1	4.2	6	13.3	9	6.5	16	7.7
	1–4 years	1	2.6	12	9.0	39	19.7	52	14.1
	5–17 years	0	0	4	1.2	40	15.0	44	7.0
Sex	Female	0	0	15	7.2	48	12.7	63	9.8
	Male	4	3.7	19	4.8	70	14.3	93	9.3
Jurisdiction ^d	ACT	0	0	2	10.0	1	4.5	3	6.7
	NSW	0	0	15	8.7	40	13.8	55	10.4
	NT	0	0	4	13.8	2	4.9	6	8.3
	Qld	0	0	1	1.1	9	7.8	10	4.7
	SA	1	7.7	3	5.3	8	12.7	12	9.0
	Tas.	0	0	0	0	2	10.0	2	5.1
	Vic.	3	5.4	5	3.3	50	20.9	58	13.0
	WA	0	0	4	5.5	6	7.9	10	6.3
	Median	38.5	—	14.5	—	13	—	14	—
	IQR ^c	21.0–56.5	—	10.0–19.8	—	8.3–23.5	—	9.0–24.5	—
30-day outcome	Died	0	0	1	16.7	8	19.0	9	16.7
	Survived	4	2.7	28	5.4	98	14.0	130	9.5
	Unknown	0	0	5	5.9	12	9.8	17	7.8
Onset	Community	0	0	18	3.8	58	9.9	76	6.7
	Hospital	4	4.1	16	12.4	60	21.1	80	15.7
Device-related infection	Yes	2	2.8	7	6.5	44	20.8	53	13.5
	No	2	2.2	23	4.9	65	10.8	90	7.7
	Unknown	0	0	4	16.0	9	16.4	13	15.3
Polymicrobial		3	5.0	4	9.3	11	11.0	18	8.9

a n: number of MDR isolates; %: proportion of MDR isolates compared to all tested isolates of the indicated organism.

b The numbers in each organism group may not add up to the total column, as some patients have polymicrobial bacteraemic episodes and are reported in each respective column to their episode, but only counted once in the total number of patients.

c IQR: interquartile range.

d ACT: Australian Capital Territory; NSW: New South Wales; NT: Northern Territory; Qld: Queensland; SA: South Australia; Tas.: Tasmania; Vic.: Victoria; WA: Western Australia.

Only three *P. aeruginosa* were identified as MDR (3/61; 4.9%) – one from New South Wales and two from South Australia.

Forty-one *Acinetobacter* spp. isolates were reported to AGAR: most were *A. baumannii* complex (18/41; 43.9%). All *Acinetobacter* isolates were meropenem susceptible. One *Acinetobacter* isolate was reported as ciprofloxacin resistant and one isolate was resistant to amikacin. Three isolates were resistant to co-trimoxazole (8.6%; 95%CI: 1.8–23.1%), and all resistant isolates were reported from patients aged 1–4 years.

Staphylococcus aureus

There were 607 *S. aureus* isolates reported to AGAR: 303 in 2020 and 304 in 2021. Overall, 12.9% of *S. aureus* were methicillin resistant (n: 78) and 5.6% of isolates were MDR (n: 34). The median age was 6 years, and the number of episodes reported across different age groups was similar. The majority of isolates were reported from New South Wales and Victoria. Most episodes were community-onset and were monomicrobial events (Table 1).

The Northern Territory reported the highest proportion of *S. aureus* isolates that were MRSA (45%). MRSA was not reported in the Australian Capital Territory or Tasmania. MRSA isolates were most frequently reported in patients aged 1–4 years and in hospital-onset infections (Table 4).

Overall, 12.4% of *S. aureus* were clindamycin resistant (n: 75; 95% CI: 9.8–15.2). The Northern Territory had the highest proportion of clindamycin-resistant isolates. No clindamycin-resistant isolates were reported in Tasmania. All age groups < 5 years of age reported approximately 15% of isolates resistant to clindamycin (Table 4).

Five percent of *S. aureus* were ciprofloxacin resistant (n: 32; 5.3%; 95% CI: 3.6–7.4%); 3.6% of *S. aureus* isolates were reported as resistant in 2020 (n: 11), and 6.9% in 2021 (n: 21). The highest proportion of isolates reported as ciprofloxacin resistance was in New South Wales (9.8%, n: 17). No isolates were ciprofloxacin resistant in Tasmania. Resistant isolates were most frequently reported in patients aged 91–364 days (Table 4).

Overall, 13.2% of *S. aureus* were erythromycin resistant (n: 80; 95% CI: 10.6–16.1%): The proportion of resistant isolates was 11.2% (n: 34) in 2020 and 15.1% (n: 46) in 2021. The highest proportion of erythromycin resistant isolates was reported in the Northern Territory, and from neonatal patients (Table 4).

No *S. aureus* isolates were reported trimethoprim-sulfamethoxazole or vancomycin resistant.

Overall, 5.6% of all *S. aureus* were MDR (n: 34), 64.7% of which were MRSA (n: 22). A statistically significant difference in age was identified in patients with a MDR *S. aureus* episode compared to patients with a non-MDR *S. aureus* ($p < 0.001$). MDR *S. aureus* episodes are more likely to be hospital-onset ($p < 0.001$); however, MDR infections were not more device-related ($p = 0.06$). The largest number of all MDR *S. aureus* isolates were from New South Wales, whilst the Northern Territory and the Australian Capital Territory had the highest proportion of *S. aureus* that were MDR (Table 3).

Enterococcus spp.

Overall, 170 enterococci were reported to AGAR, 95 isolates in 2020 and 75 isolates in 2021. Of the 170 isolates, 122 were *E. faecalis* and 41 were *E. faecium*. Five *E. faecium* isolates were MDR. Enterococci isolates were most frequently reported from patients < 1 year old. The largest proportions of episodes were from New South Wales and Victoria. The proportion of patients still in hospital at 30 days was higher in patients who had an enterococcal infection (Table 1).

Overall, 19.6% of isolates were ampicillin resistant (n: 33; 95% CI: 13.9–26.5%); one isolate was *E. faecalis*, whilst the rest were *E. faecium* (n: 32). Over 70% of ampicillin-resistant isolates were hospital-onset. Most isolates were reported in patients aged 5–17 years, and from patients living in New South Wales. Enterococci from hospital-onset episodes were more frequently ampicillin resistant compared to community-onset enterococcal episodes (Table 5).

Fewer than five percent of Enterococci isolates were vancomycin resistant (4.7%; 95% CI: 2.1–9.1%). Eight vancomycin-resistant *E. faecium* (VREfm) were identified (19.5%; 95% CI: 8.8–34.9%), all from hospital-onset episodes. VREfm isolates were identified in Queensland, Victoria, and New South Wales. Three *E. faecium* were identified as teicoplanin resistant (7.3%; 95% CI: 1.5–19.9%); all were from hospital-onset episodes. Teicoplanin resistant isolates were identified in New South Wales and Victoria (Table 5).

Five Enterococci were MDR (2.9%), all *E. faecium*. Four of the five MDR *Enterococcus* were identified in Victoria; the remaining isolate was identified in South Australia (Table 3).

Table 4: Number of *Staphylococcus aureus* isolates tested and proportion of isolates resistant to clindamycin, ciprofloxacin, and erythromycin from children aged < 18 years as reported to AGAR, January 2020 – December 2021

Category	Characteristic	N ^a	Clindamycin ^b			Ciprofloxacin ^b			Erythromycin ^b			
			n	R%	95% CI	n	R%	95% CI	n	R%	95% CI	
Overall		607	75	12.4	9.8–15.2	32	5.3	3.6–7.4	80	13.2	10.6–16.1	
Jurisdiction ^c	ACT	20	3	15	3.2–37.9	1	5.0	0.1–24.9	2	10	1.7–29.3	
	NSW	173	29	16.8	11.5–23.2	17	9.8	5.8–15.3	28	16.2	11.3–22.2	
	NT	29	6	20.7	8.0–39.7	1	3.4	0.1–17.8	6	20.7	8.8–38.2	
	Qld	88	13	14.8	8.1–23.9	4	4.5	1.3–11.2	14	15.9	9.4–24.7	
	SA	57	3	5.3	1.1–14.6	3	5.3	1.1–14.6	7	12.3	5.5–22.8	
	Tas.	15	0	0.0	—	0	0.0	—	0	0.0	—	
	Vic.	152	15	9.9	5.6–15.8	4	2.6	0.7–6.6	17	11.2	6.9–17.0	
	WA	73	6	8.2	3.1–17.0	2	2.7	0.3–9.5	6	8.2	3.4–16.3	
	Age group	≤ 28 days	51	9	17.6	8.4–30.9	4	7.8	2.2–18.9	10	19.6	9.8–33.1
		29–90 days	51	8	15.7	7.0–28.6	2	3.9	0.5–13.5	9	17.6	8.4–30.9
91–364 days		46	7	15.2	6.3–28.9	6	13	4.9–26.3	5	10.9	3.6–23.6	
1–4 years		133	21	15.8	1.0–23.1	8	6.0	2.6–11.5	24	18	11.9–25.6	
5–17 years		326	30	9.2	6.3–12.9	12	3.7	1.9–6.3	32	9.8	6.8–13.6	
Onset	Community	478	55	11.5	8.8–4.7	22	4.6	2.9–6.9	59	12.3	9.5–15.6	
	Hospital	129	20	15.5	9.7–22.9	10	7.8	3.8–13.8	21	16.3	10.4–23.8	
Methicillin resistance ^d	MRSA	78	15	19.2	11.2–29.7	13	16.7	9.2–26.8	18	23.1	14.3–34.0	
	MSSA	529	60	11.3	8.8–14.4	19	3.6	2.2–5.6	62	11.7	9.1–14.8	

a N: number of isolates tested.

b n: number of resistant isolates; R%: proportion of resistant isolates; 95% CI: 95% confidence interval.

c ACT: Australian Capital Territory; NSW: New South Wales; NT: Northern Territory; Qld: Queensland; SA: South Australia; Tas.: Tasmania; Vic.: Victoria; WA: Western Australia.

d MRSA: methicillin-resistant *S. aureus*; MSSA: methicillin-susceptible *S. aureus*.

Table 5: Number of *Enterococcus* spp. isolates tested and proportion of isolates resistant to ampicillin, vancomycin, and teicoplanin from children aged < 18 years as reported to AGAR, January 2020 – December 2021

Category	Characteristic	Ampicillin ^a			Vancomycin ^a			Teicoplanin ^a					
		N	n	R%	95% CI	N	n	R%	95% CI	N	n	R%	95% CI
Overall		168	33	19.6	13.9–26.5	170	8	4.7	2.1–9.1	170	3	1.8	0.4–5.1
Jurisdiction ^b	ACT	3	1	33.3	—	3	0	0	—	3	0	0	—
	NSW	68	15	22.7	13.3–34.7	68	2	2.9	0.4–10.2	68	2	2.9	0.4–10.2
	NT	2	1	50	—	2	0	0	—	2	0	0	—
	Qld	10	0	0	—	10	0	0	—	10	0	0	—
	SA	14	7	50	23.0–77.0	14	2	14.3	1.8–42.8	14	0	0	—
	Tas.	4	1	25	—	4	0	0	—	4	0	0	—
	Vic.	58	8	13.8	6.1–25.4	58	4	6.9	1.9–16.7	58	1	1.7	0.0–9.2
WA	11	0	0	—	11	0	0	—	11	0	0	—	
Age group	≤ 28 days	37	2	5.4	0.7–18.2	38	1	2.6	0.1–13.8	38	0	0	—
	29–90 days	26	2	7.7	0.9–25.1	26	1	3.8	0.1–19.6	26	0	0	—
	91–364 days	23	3	13	2.8–33.6	24	1	4.2	0.1–21.1	24	0	0	—
	1–4 years	41	7	17.1	7.2–32.1	41	3	7.3	1.5–19.9	41	2	4.9	0.6–16.5
	5–17 years	41	19	46.3	30.7–62.6	41	2	4.9	0.6–16.5	41	1	2.4	0.1–12.9
Onset	Community	71	9	12.7	6.0–22.7	71	0	0	—	72	0	0	—
	Hospital	97	24	24.7	16.5–34.5	98	8	8.2	3.6–15.5	98	3	3.1	0.6–8.7

a N: number of isolates tested; n: number of resistant isolates; R%: proportion of resistant isolates; 95% CI: 95% confidence interval.

b ACT: Australian Capital Territory; NSW: New South Wales; NT: Northern Territory; Qld: Queensland; SA: South Australia; Tas.: Tasmania; Vic.: Victoria; WA: Western Australia.

Discussion

This AGAR Kids report is the first comprehensive AGAR report describing the epidemiology of AMR in the Australian paediatric population. The AGAR surveillance programs provide the opportunity to specifically analyse and report on bacteraemia from Australian children. The programs also provide an opportunity to compare paediatric data with the Australian adult population. Although there are some similarities, numerous differences warrant ongoing paediatric specific reporting.

AMR surveillance of paediatric bacteraemia is a powerful public health tool, providing an understanding of the aetiology of disease, susceptibility patterns, and the impact of infection prevention and control strategies.^{3,10} AMR surveillance can also point to potential targets for therapeutics specifically for paediatrics: the development of medicines for children lags unacceptably behind that for adults, and industry and funders remain unguided by robust health costing data to prioritise paediatric AMR strategies.¹¹ It is estimated that every year, globally, AMR infection result in 700,000 deaths, including 200,000 newborns.¹²

There are few reports on paediatric populations from similar AMR national surveillance systems. Whilst the European system (EARS-Net) requests age to the nearest year, the data point is not mandatory and thus EARS-Net data are not routinely stratified by age.² In the UK, the English Surveillance Programme for Antimicrobial Utilisation and Resistance (ESPAUR) does stratify patient age, but paediatric-specific reports are not available.¹³ The Canadian AMR surveillance program presents key highlights of age-stratified data.¹⁴ One-off epidemiological reports have been produced from different regions and countries, yet interpretation and comparison to AGAR Kids is difficult because of differences in the organisms selected for surveillance.^{2,4,10,15–23}

This report is one of the most detailed paediatric-specific reports to date and facilitates the national public health benchmarking of AMR in children and neonates to ensure the effectiveness of AMR programs. In Australian children (excluding neonates), the most frequently reported organism was *S. aureus* (36% of isolates) and *E. coli* (23%); in neonates and the adult population this is reversed, with *E. coli* being the most frequently reported organism (43.1% and 39% respectively) followed by *S. aureus* (20.2% and 20.9% respectively). A similar observation has been identified in Japan by Kusama et al.,²² and likely reflects the commonality of urinary tract infections as the predominant source of bacteraemia in the neonatal and adult population and bone and joint infection the predominant source in children.²⁴ Several papers globally have demonstrated that *E. coli* bacteraemia is more frequent in children < 1 year of age and *S. aureus* is more frequent in children with increasing age.^{10,16,19,25–28}

In enterococcal bacteraemia, differences in the predominant species in adults and children were identified. In paediatrics, whilst approximately 25% of enterococci were identified as *E. faecium*, in adults almost 40% of episodes were due to *E. faecium*. Similarly, in several studies assessing paediatric bacteraemia, *E. faecalis* was more frequently identified than *E. faecium*.^{15,29} Interestingly, on a population level, the UK has reported a shift in enterococcal bacteraemia, with more *E. faecium* and less *E. faecalis* reported during the coronavirus disease 2019 (COVID-19) pandemic; future AGAR Kids reports will be needed to investigate if a similar trend occurs in the Australian paediatric population.¹³ Additionally, whilst when compared to the EARS-Net data, Australia ranks in the top third of countries for vancomycin-resistant *E. faecium* (VREfm), the proportion of resistance in children is significantly lower: in 2021, AGAR reported 37.9% VREfm, whilst in children there were only eight VREfm isolates (19.5%). This vast difference is important in the way infection prevention and stewardship programs need to be targeted for specific populations.

Regional differences were noted in the pathogens causing paediatric bacteraemia. The increased proportion of *Salmonella* bacteraemia in Queensland is consistent with previous Australian reports,^{17,26} and is consistent with the literature of bacteraemia in tropical environments.^{30–32} Significant differences in the antimicrobial susceptibility patterns were also observed between jurisdictions. The proportions of Enterobacterales in Victoria resistant to gentamicin/tobramycin and to third-generation cephalosporins were both significantly higher than the national average: 16.8% compared to 11.6% for gentamicin/tobramycin (p : 0.04) and 19.3% compared to 12.9% for third-generation cephalosporins (p : 0.01). Similar to gram-negative bacteraemia, regional differences in *S. aureus* were noted; there was a clear disproportion in the geographic distribution of MRSA episodes reported across Australia, and the over representation from the NT is found in both adults and children. The higher proportions of resistance in MRSA compared to MSSA and an older age group is aligned with the literature.¹³ A more detailed investigation in the geographic distribution of pathogens in paediatrics is proposed and it is hypothesised that there will be a higher proportion of MRSA across the north of Australia which has been reported in the literature.^{33–36} These geographical differences have important implications for jurisdictional infection prevention and stewardship programs.

As per the WHO-defined priority list of pathogens,³⁷ only four critical priority isolates were identified: two *P. aeruginosa* and two *E. cloacae* complex carbapenem-resistant (CRE). However, 21 WHO-defined high priority pathogens were identified other than MRSA: eight vancomycin-resistant *E. faecium* and 13 fluoroquinolone-resistant *Salmonella*. These findings suggest that despite the increasing levels of AMR in the general Australian population, WHO priority pathogens are rarely detected in the Australian paediatric population. This also demonstrates the benefit of using local paediatric data to strengthen antimicrobial stewardship programs and inform antibiotic guidelines.

The AGAR surveillance system only captures *S. aureus*, *Enterococcus* spp., Enterobacterales, *P. aeruginosa* and *Acinetobacter* spp. This limits the understanding of the true burden of bacteraemia, as key pathogens such as *Streptococcus* spp. are not recorded. Additionally, vaccine-preventable bacteraemic pathogens are not reported, and thus any changes in incidence or proportion of aetiology of bacteraemia is not captured.

Additional data analysis will be undertaken on a regular basis to track the emergence of AMR in Australian children, to investigate the differences in AMR between adults and paediatric bacteraemia isolates, and to assess the trends in paediatric AMR across Australia.

Overall, this report highlights key differences in antimicrobial resistance and geographic distributions of bacteraemic pathogens in Australian children. These differences emphasise the importance of paediatric-specific infection prevention and stewardship programs, and the need to continue AMR surveillance in the paediatric population.

Acknowledgments

This report is from the AGAR-Kids group and funded by the Australian Government Department of Health and Aged Care.

Members of AGAR in 2020/2021 were:

Australian Capital Territory

Peter Collignon and Susan Bradbury, The Canberra Hospital

New South Wales

Thomas Gottlieb and John Huynh, Concord Hospital

James Branley and Linda Douglass, Nepean Hospital

Angela Wong, Royal North Shore Hospital

Jon Iredell and Andrew Ginn, Westmead Hospital

Rod Givney and Bree Harris, John Hunter Hospital

Peter Newton and Melissa Hoddle, Wollongong Hospital

David Lorenz, St Vincent's Hospital

Monica Lahra and Peter Huntington, Sydney Children's Hospital

Alison Kesson and Andrew Jarrett, Children's Hospital Westmead

Michael Maley and Helen Ziochos, Liverpool Hospital

Northern Territory

Sebastian van Hal and Alicia Beukers, Royal Prince Alfred Hospital Northern Territory

Rob Baird and Jann Hennessy, Royal Darwin Hospital

James McLeod, Alice Springs Hospital

Queensland

Graeme Nimmo and Narelle George, Pathology Queensland Central Laboratory, Royal Brisbane and Women's Hospital

Petra Derrington and Cheryl Curtis, Pathology Queensland Gold Coast Hospital

Robert Horvath and Laura Martin, Pathology Queensland Prince Charles Hospital

Naomi Runnegar and Joel Douglas, Pathology Queensland Princess Alexandra Hospital

Jennifer Robson and Marianne Allen, Sullivan Nicolaides Pathology, Greenslopes Hospital

Clare Nourse, Queensland Children's Hospital

South Australia

Kelly Papanou and Xiao Ming Chen, SA Pathology (Flinders Medical Centre)

Morgyn Warner and Kija Smith, SA Pathology (Royal Adelaide Hospital and Women's and Children's Hospital)

Tasmania

Louise Cooley and David Jones, Royal Hobart Hospital

Pankaja Kalukottege, Kathy Wilcox and Brooke Wooley, Launceston General Hospital

Victoria

Denis Spelman, Adam Jenney and Jacqueline Williams, The Alfred Hospital

Marcel Leroi and Elizabeth Grabsch, Austin Health

Tony Korman, Despina Kotsanas and Kathryn Cisera, Monash Medical Centre, Monash Children's Hospital and Dandenong Hospital

Andrew Daley and Gena Gonis, Royal Women's and Children's Hospital

Mary Jo Waters, Amy Crowe and Lisa Brenton, St Vincent's Hospital

Katherine Bond and Rose Cotronei, Royal Melbourne Hospital

Western Australia

David McGeachie and Denise Daley, PathWest Laboratory Medicine – WA Fiona Stanley Hospital

Ronan Murray and Jacinta Bowman, PathWest Laboratory Medicine – WA Sir Charles Gairdner Hospital

Michael Leung, PathWest Laboratory Medicine – Northwest WA

Owen Robinson and Geoffrey Coombs, PathWest Laboratory Medicine – WA Royal Perth Hospital

Sudha Pottumarthy-Boddu and Jacqueline Foster, Australian Clinical Laboratories, St John of God Hospital, Murdoch

Shalinie Perera and Ian Meyer, Western Diagnostic Pathology, Joondalup Hospital

Christopher Blyth, Perth Children's Hospital

Author details

Anita Williams,¹

Geoffrey W Coombs,^{2,3,4}

Jan Bell,⁵

Denise A Daley,^{3,4}

Shakeel Mowlaboccus,^{2,3,4}

Penelope A Bryant,^{6,7}

Anita J Campbell,^{1,9,10}

Louise Cooley,¹¹

Jon Iredell,^{12,13,14,15}

Adam D Irwin,^{16,17}

Alison Kesson,^{13,18}

Brendan McMullan,^{19,20}

Morgyn S Warner,^{21,22}

Phoebe Williams,^{13,20,23}

Christopher C Blyth.^{1,9,10,24}

1. Wesfarmers Centre of Vaccines and Infectious Diseases, Telethon Kids Institute, Nedlands, WA
2. Antimicrobial Resistance and Infectious Diseases (AMRID) Research Laboratory, Murdoch University, Murdoch, WA
3. Department of Microbiology, PathWest Laboratory Medicine WA, Fiona Stanley Hospital, Murdoch, WA
4. Australian Group on Antimicrobial Resistance, Fiona Stanley Hospital, Murdoch, WA
5. Australian Group on Antimicrobial Resistance, Adelaide, SA
6. Infectious Diseases, The Royal Children's Hospital Melbourne, Parkville, VIC
7. Department of Paediatrics, University of Melbourne, Parkville, Victoria
8. Murdoch Children's Research Institute, Parkville, Victoria
9. Department of Infectious Diseases, Perth Children's Hospital, Nedlands, WA
10. Division of Paediatrics, School of Medicine, University of Western Australia, Nedlands, WA
11. Department of Infectious Diseases, Royal Hobart Hospital, Tasmania

12. Centre for Infectious Diseases and Microbiology, The Westmead Institute for Medical Research, Westmead, NSW
13. Faculty of Medicine and Health, University of Sydney, Sydney, NSW
14. Westmead Hospital, Western Sydney Local Health District, Westmead, NSW
15. Sydney Institute for Infectious Disease, The University of Sydney, Sydney, NSW
16. Centre for Clinical Research, University of Queensland, Brisbane, Queensland
17. Infection Management Prevention Service, Queensland Children's Hospital, Brisbane, Queensland
18. Department of Infectious Disease & Microbiology, The Children's Hospital at Westmead, Westmead, NSW
19. School of Clinical Medicine, Faculty of Medicine and Health, University of New South Wales, Kensington, NSW
20. Department of Immunology and Infectious Diseases, Sydney Children's Hospital, Sydney Children's Hospitals Network, Randwick, NSW
21. Infectious Diseases Unit, Central Adelaide Local Health Network, Adelaide, SA
22. Faculty of Health and Medical Sciences, University of Adelaide, Adelaide, SA
23. School of Women and Children's Health, University of New South Wales, Kensington, NSW
24. Department of Microbiology, PathWest Laboratory Medicine WA, QEII Medical Centre, Nedlands, WA

Corresponding author

Ms Anita Williams

Address: Telethon Kids Institute, 15 Hospital Ave, Nedlands 6009 WA

Phone: +61 8 6319 1000

Email: anita.williams@telethonkids.org.au

References

1. BATTERY J, YANG Y, SHARLAND M, World Society for Pediatric Infectious Diseases. World Society for Pediatric Infectious Diseases declaration on combating antimicrobial resistance in children. *World J Pediatr*. 2018;14(6):523–4. doi: <https://doi.org/10.1007/s12519-018-0195-x>.
2. Bielicki JA, Lundin R, Sharland M, ARPEC Project. Antibiotic resistance prevalence in routine bloodstream isolates from children's hospitals varies substantially from adult surveillance data in Europe. *Pediatr Infect Dis J*. 2015;34(7):734–41. doi: <https://doi.org/10.1097/INF.0000000000000652>.
3. Pai S, Enoch DA, Aliyu SH. Bacteremia in children: epidemiology, clinical diagnosis and antibiotic treatment. *Expert Rev Anti Infect Ther*. 2015;13(9):1073–88. doi: <https://doi.org/10.1586/14787210.2015.1063418>.
4. Campbell AJ, Daley DA, Bell JM, Pang S, Coombs GW, Carapetis JR et al. Progress towards a coordinated, national paediatric antimicrobial resistance surveillance programme: *Staphylococcus aureus*, enterococcal and gram-negative bacteraemia in Australia. *J Antimicrob Chemother*. 2020;75(6):1639–44. doi: <https://doi.org/10.1093/jac/dkaa065>.
5. Murdoch F, Danial J, Morris AK, Czarniak E, Bishop JL, Glass E et al. The Scottish enhanced *Staphylococcus aureus* bacteraemia surveillance programme: the first 18 months of data in children. *J Hosp Infect*. 2017;97(2):127–32. doi: <https://doi.org/10.1016/j.jhin.2017.06.017>.
6. Australian Institute of Health and Welfare (AIHW). *Australian Hospital Peer Groups. Health Services Series No. 66*. Canberra: AIHW; 2015. Available from: <https://www.aihw.gov.au/getmedia/79e7d756-7cfe-49bf-b8c0-0bbb0daa2430/14825.pdf>.
7. The European Committee on Antimicrobial Susceptibility Testing (EUCAST). *Breakpoint tables for interpretation of MICs and zone diameters. Version 12.0*. Basel: EUCAST; 1 January 2022. Available from: https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_12.0_Breakpoint_Tables.xlsx.
8. Berends MS, Friedrich AW, Albers CJ, Luz CF, Sinha BNM, Glasner C. AMR – an R package for working with antimicrobial resistance data. *J Stat Softw*. 2022;104(3):1–31. doi: <https://doi.org/10.18637/jss.v104.i03>.
9. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect*. 2012;18(3):268–81. doi: <https://doi.org/10.1111/j.1469-0691.2011.03570.x>.
10. Henderson KL, Johnson AP, Muller-Pebody B, Charlett A, Gilbert R, Sharland M. The changing aetiology of paediatric bacteraemia in England and Wales, 1998–2007. *J Med Microbiol*. 2010;59(2):213–19. doi: <https://doi.org/10.1099/jmm.0.015271-0>.
11. Pana ZD, El-Shabrawi M, Sultan MA, Murray T, Alam A, Yewale V et al. Fighting the hidden pandemic of antimicrobial resistance in paediatrics: recommendations from the International Pediatric Association. *BMJ Paediatr Open*. 2023;7(1):e002084. doi: <https://doi.org/10.1136/bmjpo-2023-002084>.
12. Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet*. 2022;399(10325):629–55. doi: [https://doi.org/10.1016/S0140-6736\(21\)02724-0](https://doi.org/10.1016/S0140-6736(21)02724-0).
13. United Kingdom Health Security Agency. *English surveillance programme for antimicrobial utilisation and resistance (ESPAUR) report 2021 to 2022*. London: United Kingdom Health Security Agency; 21 November 2022. Available from: <https://webarchive.nationalarchives.gov.uk/ukgwa/20231002172235/https://www.gov.uk/government/publications/english-surveillance-programme-antimicrobial-utilisation-and-resistance-espaur-report>.

14. Antimicrobial Resistance Taskforce (AMRTF). *Système canadien de surveillance de la résistance aux antimicrobiens : Rapport de 2022*. Ottawa, Canada: Public Health Agency of Canada (PHAC); 28 November 2022. doi: <https://doi.org/10.58333/f241022>.
15. Bizot E, Truong J, Mariani-Kurkdjian P, Bonacorsi S, Faye A, Caseris M. Pediatric enterococcal bacteremia: a 12-year retrospective study in a French pediatric center. *Pediatr Infect Dis J*. 2022;41(9):e346–50. doi: <https://doi.org/10.1097/INF.0000000000003631>.
16. Buetti N, Atkinson A, Kottanattu L, Bielicki J, Marschall J, Kronenberg A. Patterns and trends of pediatric bloodstream infections: a 7-year surveillance study. *Eur J Clin Microbiol Infect Dis*. 2017;36(3):537–44. doi: <https://doi.org/10.1007/s10096-016-2830-6>.
17. Douglas MW, Lum G, Roy J, Fisher DA, Anstey NM, Currie BJ. Epidemiology of community-acquired and nosocomial bloodstream infections in tropical Australia: a 12-month prospective study. *Trop Med Int Health*. 2004;9(7):795–804. doi: <https://doi.org/10.1111/j.1365-3156.2004.01269.x>.
18. Droz N, Hsia Y, Ellis S, Dramowski A, Sharland M, Basmaci R. Bacterial pathogens and resistance causing community acquired paediatric bloodstream infections in low- and middle-income countries: a systematic review and meta-analysis. *Antimicrob Resist Infect Control*. 2019;8(1):207. doi: <https://doi.org/10.1186/s13756-019-0673-5>.
19. Ferreira M, Santos M, Rodrigues J, Diogo C, Resende C, Baptista C et al. Epidemiology of bacteremia in a pediatric population – a 10-year study. *Enferm Infecc Microbiol Clin (Engl Ed)*. 2021:S0213-005X(21)00207-X. doi: <https://doi.org/10.1016/j.eimc.2021.06.011>.
20. Graff KE, Windsor WJ, Calvimontes DM, Melgar MA, Galvez N, Rivera JG et al. Antimicrobial resistance trends at a pediatric hospital in Guatemala City, 2005–2019. *J Pediatric Infect Dis Soc*. 2021:piab048. doi: <https://doi.org/10.1093/jpids/piab048>.
21. Iroh Tam P-Y, Musicha P, Kawaza K, Cornick J, Denis B, Freyne B et al. Emerging resistance to empiric antimicrobial regimens for pediatric bloodstream infections in Malawi (1998–2017). *Clin Infect Dis*. 2019;69(1):61–8. doi: <https://doi.org/10.1093/cid/ciy834>.
22. Kusama Y, Ito K, Fukuda H, Matsunaga N, Ohmagari N. National database study of trends in bacteraemia aetiology among children and adults in Japan: a longitudinal observational study. *BMJ Open*. 2021;11(3):e043774. doi: <https://doi.org/10.1136/bmjopen-2020-043774>.
23. Agyeman PKA, Schlapbach LJ, Giannoni E, Stocker M, Posfay-Barbe KM, Heininger U et al. Epidemiology of blood culture-proven bacterial sepsis in children in Switzerland: a population-based cohort study. *Lancet Child Adolesc Health*. 2017;1(2):124–33. doi: [https://doi.org/10.1016/S2352-4642\(17\)30010-X](https://doi.org/10.1016/S2352-4642(17)30010-X).
24. Coombs G, Bell JM, Daley D, Collignon P, Cooley L, Gottlieb T et al. *Australian Group on Antimicrobial Resistance (AGAR) Surveillance Outcomes Programs: Bloodstream infections 2021 Report*. Sydney: Australian Commission on Safety and Quality in Health Care; November 2022. Available from: https://www.safetyandquality.gov.au/sites/default/files/2022-12/2021_agar_amalgam_report.pdf.
25. Laupland KB, Gregson DB, Vanderkooi OG, Ross T, Kellner JD. The changing burden of pediatric bloodstream infections in Calgary, Canada, 2000–2006. *Pediatr Infect Dis J*. 2009;28(2):114–7. doi: <https://doi.org/10.1097/INF.0b013e318187ad5a>.
26. Er J, Wallis P, Maloney S, Norton R. Paediatric bacteraemias in tropical Australia. *J Paediatr Child Health*. 2015;51(4):437–42. doi: <https://doi.org/10.1111/jpc.12750>.
27. Shakur SM, Whitehall J, Mudgil P. Pediatric bloodstream infections in metropolitan Australia. *World J Pediatr*. 2019;15(2):161–7. doi: <https://doi.org/10.1007/s12519-018-00221-3>.

28. Qiu Y, Yang J, Chen Y, Yang J, Zhu Q, Zhu C et al. Microbiological profiles and antimicrobial resistance patterns of pediatric bloodstream pathogens in China, 2016–2018. *Eur J Clin Microbiol Infect Dis*. 2021;40(4):739–49. doi: <https://doi.org/10.1007/s10096-020-04069-2>.
29. Stover KR, Morrison A, Collier T, Schneider E, Wagner JL, Capino AC et al. Epidemiology and risk factors for bacteremia in pediatric and adolescent patients. *J Pharm Pract*. 2021;34(3):360–4. doi: <https://doi.org/10.1177/0897190019868056>.
30. Chang YJ, Chen YC, Chen NW, Hsu YJ, Chu HH, Chen CL et al. Changing antimicrobial resistance and epidemiology of non-typhoidal *Salmonella* infection in Taiwanese children. *Front Microbiol*. 2021;12:648008. doi: <https://doi.org/10.3389/fmicb.2021.648008>.
31. Moore N, Ashley EA, Dickson B, Douangnouvong A, Panyaviseth P, Turner P et al. A systematic review of antimicrobial resistance among children in low- and middle-income countries in the Western Pacific Region (WPRO). Amsterdam: Elsevier, Social Sciences Research Network (SSRN); 4 May 2023. [Accessed on 23 June 2023.] doi: <https://doi.org/10.2139/ssrn.4430748>.
32. Parisi A, Crump JA, Stafford R, Glass K, Howden BP, Kirk MD. Increasing incidence of invasive nontyphoidal *Salmonella* infections in Queensland, Australia, 2007–2016. *PLoS Negl Trop Dis*. 2019;13(3):e0007187. doi: <https://doi.org/10.1371/journal.pntd.0007187>.
33. McMullan BJ, Bowen A, Blyth CC, Van Hal S, Korman TM, BATTERY J et al. Epidemiology and mortality of *Staphylococcus aureus* bacteremia in Australian and New Zealand children. *JAMA Pediatr*. 2016;170(10):979–86. doi: <https://doi.org/10.1001/jamapediatrics.2016.1477>.
34. Harch SAJ, MacMorran E, Tong SYC, Holt DC, Wilson J, Athan E et al. High burden of complicated skin and soft tissue infections in the Indigenous population of Central Australia due to dominant Pantone Valentine leucocidin clones ST93-MRSA and CC121-MSSA. *BMC Infect Dis*. 2017;17(1):405. doi: <https://doi.org/10.1186/s12879-017-2460-3>.
35. Engelman D, Hofer A, Davis JS, Carapetis JR, Baird RW, Giffard PM et al. Invasive *Staphylococcus aureus* infections in children in tropical northern Australia. *J Pediatr Infect Dis Soc*. 2014;3(4):304–11. doi: <https://doi.org/10.1093/jpids/piu013>.
36. Hewagama S, Spelman T, Woolley M, McLeod J, Gordon D, Einsiedel L. The epidemiology of *Staphylococcus aureus* and Pantone-Valentine leucocidin (pvl) in Central Australia, 2006–2010. *BMC Infect Dis*. 2016;16:382. doi: <https://doi.org/10.1186/s12879-016-1698-5>.
37. World Health Organization (WHO). *WHO Bacterial Priority Pathogens List, 2024: Bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance*. Geneva: WHO; 17 May 2024. Available from: <https://www.who.int/publications/i/item/9789240093461>.