

Letter to the Editor

Maternal vaginal colonisation with *Neisseria meningitidis* serogroup B and late onset neonatal invasive meningococcal disease

Nadiya Brell, Chloe Story, Sebastiaan J van Hal, Martin A Weber,
Monica M Lahra

Further to a recent report of an outbreak of *Neisseria meningitidis* serogroup Y serotype (ST-) 1466 urogenital infections in Australia,¹ we report a case of late-onset neonatal invasive meningococcal disease (IMD) secondary to maternal vaginal colonisation. The risk associated with vaginal colonisation of *N. meningitidis* for vertical transmission is unknown, but is of heightened concern given the increased detection of urogenital infections and the clinical and public health implications.¹ Isolated cases of neonatal IMD associated with either maternal IMD or colonisation predating the recent outbreak have been reported previously.^{2,3}

A 21-year-old woman (G1P0, 38⁺³ weeks gestation) presented with spontaneous rupture of membranes (SROM), afebrile, with clear liquor. A vaginal swab was collected, but results were pending at the time of delivery. She did not meet clinical risk criteria for Group B *Streptococcus* (GBS) prophylaxis,⁴ and did not receive antibiotics in labour. She proceeded to emergency caesarean section for foetal distress five hours after SROM. The 3,195 g neonate had APGAR scores of 5 and 6 and was treated empirically with intravenous ampicillin and gentamicin for 48 hours. The neonate had no source of sepsis identified, including a set of negative blood cultures collected prior to antibiotics. They were discharged home well on day four of life.

On day nine, the neonate was readmitted with fevers. At this time, it was noted that the mother's vaginal swab collected at delivery grew *N. meningitidis* and GBS. Neonatal cerebrospinal fluid grew *N. meningitidis* serogroup B; blood cultures were negative. The neonate received 14 days of antibiotics and was discharged home without evident disability.

Histological examination of the placenta showed evidence of intraamniotic infection with chorioamnionitis, umbilical panvasculitis and funisitis. *N. meningitidis* DNA was detected in placental tissue. Whole genome sequencing of neonatal and maternal isolates found both were Men-B ST-2506 (clonal complex 32) and were identical (1 SNP difference on split kmer analysis).⁵

Late-onset neonatal IMD associated with maternal vaginal *N. meningitidis* has been documented.^{2,3} This mother presented with SROM, had histological evidence of ascending intrauterine infection and *N. meningitidis* DNA detected in the placenta. Delayed onset of neonatal sepsis was likely attributable to partial treatment. This case further highlights the need to establish clear risk management strategies for vaginal *N. meningitidis* colonisation in pregnancy for mothers, neonates and their contacts.

Acknowledgments

We thank the patient for their consent to publication of this case description.

Author details

Nadiya Brell,^{1,2}

Chloe Story,¹

Sebastiaan J van Hal,^{3,4}

Martin A. Weber,^{2,5,6}

Monica M Lahra,^{2,7}

1. NSW Health Pathology, Microbiology, The Wollongong Hospital, Wollongong, New South Wales, Australia
2. The Faculty of Medicine, The University of New South Wales, Sydney, New South Wales Australia
3. NSW Health Pathology, Microbiology, Royal Prince Alfred Hospital, Sydney, New South Wales, Australia
4. Central Clinical School, University of Sydney, Sydney, New South Wales, Australia
5. NSW Health Pathology, Anatomical Pathology, The Prince of Wales Hospital, Randwick, Sydney, New South Wales, Australia
6. Faculty of Biomedical Sciences, The University of New South Wales, Sydney, New South Wales, Australia
7. World Health Organization Collaborating Centre for STI and AMR, NSW Health Pathology, Microbiology, The Prince of Wales Hospital, Randwick, Sydney, New South Wales, Australia

Corresponding author

Professor Monica M Lahra

World Health Organization Collaborating Centre for STI and AMR, NSW Health Pathology, Microbiology, The Prince of Wales Hospital, Randwick, Sydney, New South Wales, Australia

Phone: +61 2 9382 3586

Email: monica.lahra@health.nsw.gov.au

References

1. Lahra MM, Latham NH, Templeton DJ, Read P, Carmody C, Ryder N et al. Investigation and response to an outbreak of *Neisseria meningitidis* serogroup Y ST-1466 urogenital infections, Australia. *Commun Dis Intell* (2018). 2024;48. doi: <https://doi.org/10.33321/cdi.2024.48.20>.
2. Chacon-Cruz E, Alvelais-Palacios JA, Rodriguez-Valencia JA, Lopatynsky-Reyes EZ, Volker-Soberanes ML, Rivas-Landeros RM. Meningococcal neonatal purulent conjunctivitis/sepsis and asymptomatic carriage of *N. meningitidis* in mother's vagina and both parents' nasopharynx. *Case Rep Infect Dis*. 2017;2017:6132857. doi: <https://doi.org/10.1155/2017/6132857>.
3. Hart J, Dowse GK, Porter M, Speers DJ, Keil AD, Bew JD et al. Obstetric and neonatal invasive meningococcal disease caused by *Neisseria meningitidis* serogroup W, Western Australia, Australia. *Emerg Infect Dis*. 2024;30(2):368–71. doi: <https://doi.org/10.3201/eid3002.230639>.
4. Royal Australian and New Zealand College of Obstetricians and Gynaecologists (RANZCOG). *Maternal Group B Streptococcus in pregnancy: screening and management*. Melbourne: RANZCOG; July 2019. Available from: <https://ranzcog.edu.au/wp-content/uploads/Maternal-Group-B-Streptococcus-in-Pregnancy-Screening-Management.pdf>.
5. Harris SR. SKA: split kmer analysis toolkit for bacterial genomic epidemiology. *BioRxiv*. 2018;453142. doi: <https://doi.org/10.1101/453142>.



Communicable Diseases Intelligence (CDI) is a peer-reviewed scientific journal published by the Australian Centre for Disease Control.

The journal aims to disseminate information on the epidemiology, surveillance, prevention and control of communicable diseases of relevance to Australia and the near region.

Editor

Dr Elise Firman

Deputy Editor

Simon Petrie

Design and Production

Lisa Thompson

Editorial Advisory Board

David Durrheim, Mark Ferson, Clare Huppatz, John Kaldor, Martyn Kirk and Meru Sheel

Submit an Article

Submit your next communicable disease related article to CDI for consideration.

Guidelines for authors and details on how to submit your publication is available on our website, or by email to the CDI Editor.

Contact us

Communicable Diseases Intelligence (CDI)
Australian Centre for Disease Control
PO Box 795, Canberra ACT 2601
Australia

Website: cdc.gov.au/cdi

Email: cdi.editor@cdc.gov.au

© 2026 Commonwealth of Australia as represented by the Australian Centre for Disease Control

ISSN: 2209-6051 Online

This journal is indexed by Index Medicus and Medline.

Creative Commons Licence

This publication is licensed under a Creative Commons Attribution-Non-Commercial-NoDerivatives 4.0 International Licence (Licence). You must read and understand the Licence before using any material from this publication.

Restrictions

The Licence does not cover, and there is no permission given for, use of any of the following material found in this publication (if any):

- the Commonwealth Coat of Arms (by way of information, the terms under which the Coat of Arms may be used can be found on the Department of Prime Minister and Cabinet website);
- any logos (including the Australian Centre for Disease Control's logo) and trademarks;
- any photographs and images;
- any signatures; and
- any material belonging to third parties.

Disclaimer

Opinions expressed in *Communicable Diseases Intelligence* are those of the authors and not necessarily those of the Australian Government or the Australian Centre for Disease Control. Data may be subject to revision.

Enquiries

Enquiries regarding any other use of this publication should be addressed to the CDI Editor.