
Annual report

Australian Meningococcal Surveillance Programme annual report, 2016

Monica M Lahra, Rodney Enriquez for the National Neisseria Network

Abstract

In 2016, there were 243 laboratory-confirmed cases of invasive meningococcal disease analysed by the Australian National Neisseria Network. This number was the highest number of laboratory confirmed cases since 2008. Probable and laboratory confirmed invasive meningococcal disease (IMD) are notifiable in Australia, and there were 252 IMD cases notified to the National Notifiable Diseases Surveillance System in 2016, the highest number reported since 2010. Meningococcal serogrouping was able to be determined for 98% (237/243) of laboratory confirmed IMD cases. Serogroup B infections accounted for 87 cases (37%), the lowest number and proportion reported since inception of the Australian Meningococcal Surveillance Programme (AMSP) in 1997. In contrast, the number and proportion of serogroup W infections (44%, 107 cases) in 2016 was the highest since the AMSP began. In addition, the number and proportion of serogroup Y infections (16%, 40 cases) was also the highest recorded by the AMSP. Molecular typing results were available for 225 of the 243 IMD cases. Of the serogroup W IMD strains that were able to be genotyped, 92% (97/105) have the PorA antigen encoding gene type P1.5,2 and of these, 72% (70/97) were sequence type 11, the same type as the hypervirulent serogroup W strain that has been circulating in the UK and South America since 2009. The primary IMD age peak was observed in adults aged 45 years or more, whilst secondary disease peaks were observed in those aged less than 5 years, and in adolescents aged 15–19 years. Serogroup B infections predominated in the age groups less than 1 year and 20–24 years, whereas serogroup W infections predominated in those aged 45 years or more. For all other age groups, distribution of serogroup B and W infections was roughly equal. Of the IMD isolates tested for antimicrobial susceptibility, 6% (11/189) were resistant to penicillin, and decreased susceptibility to penicillin was observed in a further 90% (170/189) of isolates. One Men W isolate demonstrated an elevated minimum inhibitory concentration (MIC) to ceftriaxone (0.125mg/L), the highest reported in Australia. All isolates tested were susceptible to rifampicin and ciprofloxacin.

Keywords: antibiotic resistance; disease surveillance; meningococcal disease; Neisseria meningitidis

Introduction

Australia's National Neisseria Network (NNN) is an established, collaborative network of reference laboratories in each state and territory that contribute to the laboratory surveillance system of the pathogenic *Neisseria* species (*N. meningitidis* and *N. gonorrhoeae*). Since 1994 the NNN has coordinated laboratory data from the examination of *N. meningitidis* cases of invasive meningococcal disease (IMD) for the Australian Meningococcal Surveillance Programme

(AMSP).¹ The AMSP is funded by the Australian Government Department of Health. The NNN laboratories supply phenotypic and genotypic data of invasive meningococci for the AMSP. These data supplement the notification data from the National Notifiable Diseases Surveillance System (NNDSS), which includes cases of probable and laboratory confirmed IMD. The characteristics of meningococci responsible for IMD, and the associated demographic information, are important considerations for management of individual patients and their contacts; and

to inform public health responses for outbreaks or case clusters, locally and nationally. The introduction of the publicly funded conjugate serogroup C meningococcal vaccine onto the National Immunisation Program in 2003 has seen a significant and sustained reduction in the number of cases of serogroup C IMD after 2003.² However, IMD remains an issue of public health concern in Australia and continued monitoring of phenotypic and genotypic features of IMD strains is critical to monitor, plan and inform clinical management and public health interventions.

Methods

Case confirmation of invasive meningococcal disease

Case confirmation is based on isolation of *N. meningitidis*, or a positive nucleic acid amplification testing (NAAT) from a normally sterile site, defined as laboratory definitive evidence of IMD by the Communicable Diseases Network Australia (CDNA) criteria³. Information regarding the site of infection, age and sex of the patients is collated by the NNN for the AMSP.

IMD cases are categorised on the basis of the site from which *N. meningitidis* was isolated, or from which meningococcal DNA was detected (blood, joint fluid, vitreous fluid). When *N. meningitidis* is detected from both blood and cerebrospinal fluid (CSF) from the same patient, the case is classified as one of meningitis.

Phenotyping and genotyping of *Neisseria meningitidis*

Phenotyping is limited to the determination of the serogroup by detection of soluble polysaccharide antigens. Genotyping of both isolates and DNA extracts is performed by sequencing of products derived from amplification of the porin genes *porA*, *porB* and *fetA*.

Antibiotic susceptibility testing

Isolates were tested to determine their minimum inhibitory concentration (MIC) values to

antibiotics used for therapeutic and prophylactic purposes: ceftriaxone, ciprofloxacin; rifampicin. This program defines the penicillin categories as: sensitive (MIC \leq 0.03 mg/L); less sensitive (MIC 0.06–0.5 mg/L) and resistant (MIC \geq 1 mg/L).

Results

In 2016, there were 243 laboratory-confirmed cases of IMD analysed by the NNN, and 252 cases notified to the NNDSS. Thus, laboratory data were available for 96% of notified cases of IMD in Australia in 2016 (Figure 1). This number of laboratory-confirmed cases of IMD was the highest reported since 2008, with an increase of 40% from the previous year (n=174). The number of cases notified to the NNDSS was the highest reported since 2010 (n=226), with an increase of 39% from the previous year (n=182). In 2016 the peak incidence for IMD occurred in mid-spring and early summer (1 October to 31 December 2016) (Table 1). This was different to previous years where the peak incidence occurred in mid-winter and early spring.

Victoria reported the highest number of cases (76 cases) in 2016, an increase from 54 cases in 2015 and the highest number of cases reported from this state since 2005 (n=80) (Table 2). New South Wales had the second highest number of IMD cases in 2016 (69 cases) and this was the highest number of cases reported in this state since 2010 (n=76). All jurisdictions, with the exception of South Australia and the Australian Capital Territory, recorded a rise in IMD cases in 2016 compared with 2015.

Age distribution

The peak incidence of IMD in 2016, as in 2015, occurred in adults aged 45 years or more. This age group represented 35% (86/243) of IMD cases in 2016 (Table 3). Within this age group, 50 cases were in those aged 65 years or more, which was the highest number and proportion of cases for this age group reported by the AMSP. Prior to 2015, the primary peak incidence of IMD was in children less than 5 years of age, however in 2016, they represented 21% of IMD cases, the lowest proportion of cases noted by the AMSP

Figure 1: Number of invasive meningococcal disease cases reported to the National Notifiable Diseases Surveillance System compared with laboratory confirmed data from the Australian Meningococcal Surveillance Programme, Australia, 2016




Table 1: Laboratory confirmed cases of invasive meningococcal disease, Australia, 2016, by quarter

Serogroup	01 January - 31 March	01 April - 30 June	01 July - 30 September	01 October - 31 December	2016 Total
B	19	18	28	22	87
C	2	0	0	1	3
Y	4	8	10	18	40
W	14	17	36	40	107
NG	2	0	1	0	3
ND	1	0	1	1	3
Total	42	43	76	82	243

NG: non groupable.
ND: not determined.

Table 2: Number of laboratory confirmed cases of invasive meningococcal disease, Australia, 2016, by state or territory and serogroup

State or territory	Serogroup						Total
	B	C	Y	W	NG	ND	
ACT	1	0	0	1	0	0	2
NSW	25	2	15	23	3	1	69
NT	2	0	0	0	0	0	2
Qld	15	0	13	14	0	0	42
SA	22	0	0	4	0	0	26
Tas.	0	0	1	4	0	0	5
Vic.	17	1	9	47	0	2	76
WA	5	0	2	14	0	0	21
Australia	87	3	40	107	3	3	243
Proportion of all IMD cases	35.8%	1.2%	16.5%	44.0%	1.2%	1.2%	

NG: non groupable
 ND: not determined

Table 3: Laboratory-confirmed cases of invasive meningococcal disease, Australia, 2016, by age and serogroup

Serogroup	Age group									Total
	<1	1 to 4	5 to 9	10-14	15-19	20-24	25-44	45-64	65+	
B	14	11	3	1	14	18	15	6	5	87
C	0	0	0	0	1	0	1	1	0	3
Y	2	1	1	0	5	4	2	10	15	40
W	8	11	3	0	13	7	16	19	30	107
NG	0	2	0	0	0	0	1	0	0	3
ND	0	1	0	0	1	1	0	0	0	3
Total	24	26	7	1	34	30	35	36	50	243
%B of within age group	58.3	42.3	42.9	100.0	41.2	60.0	42.9	16.7	10.0	

NG: non groupable
 ND: not determined

for this age group in any year. Between 2003 and 2014, the proportion of IMD that occurred in children aged less than 5 years ranged from 28% to 36% of cases. A secondary disease peak has also been observed in previous years amongst adolescents aged 15–19 years. Of the total cases of IMD in 2016, 14% (34/243) were in those aged 15–19 years, which was less than the proportion reported in the period 2006 to 2011 (16–20%), and 2013–2015 (17–20%).

Anatomical site of samples for laboratory confirmed cases

In 2016, diagnosis was made by a positive culture in 78% (189/243) of cases and 22% (54/243) of cases were confirmed by NAAT testing alone (Table 4).

There were 45 diagnoses of meningitis based on cultures or NAAT examination of CSF either alone or with a positive blood sample. There

Table 4: Number of laboratory-confirmed cases of invasive meningococcal disease, Australia, 2016, by anatomical source and method of confirmation

Specimen type	Isolate of NM*	PCR positive**	Total
Blood	162	26	188
CSF +/- Blood	17	28	45
Other	10	0	10
Total	189	54	243

* NM: *Neisseria meningitidis*

** NAAT: nucleic acid amplification testing.

were 188 diagnoses of septicaemia based on cultures or NAAT examination from blood samples alone (Table 4). There were 8 IMD diagnoses by positive joint fluid culture, 1 IMD diagnosis by positive tissue culture, and 1 IMD diagnosis by positive abscess culture.

Serogroup data

Number and proportions of cases of serogroup B, C, Y, W invasive meningococcal disease

The serogroup was able to be determined for 237 of 243 laboratory-confirmed cases of IMD (98%) in 2016 (Tables 2 and 3). The overall decrease in IMD cases since 2002 was initially predominantly due to a reduction in the number of cases of IMD caused by serogroup C from 2003 to 2007 following the introduction of the serogroup C vaccine. After 2009, a decline in the numbers IMD cases caused by serogroup B was reported, from 194 cases in 2009 to 104 cases in 2013. In 2014, there was an increase in the numbers of IMD cases caused by serogroup B (n=129), however, in 2015 the numbers of IMD cases caused by serogroup B was similar to 2013. In the years 2006–2012 the proportion of IMD cases caused by serogroup B was 84%–88%, in 2013–2014 it was 75–80%, and in 2015 it was 64%. The number and proportion of IMD cases caused by serogroup B declined further in 2016 to the lowest number (n=87) and proportion

of total IMD (36%) reported by the AMSP. The number of IMD cases caused by serogroup C (3 cases) in 2016 was similar to the previous year 2015 (2 cases), which was the lowest total reported by the AMSP. Since 2014, the rise in the total number of IMD cases has been due to a rise in the number of cases of IMD caused by serogroup W and serogroup Y (Figure 2).


The number and proportion of cases of IMD caused by serogroup W in 2016 (107 cases, 44% of total IMD) was the highest reported by the AMSP, and was almost triple the number of cases reported in 2015 (36 cases), and a ten-fold increase in the average number of annual IMD cases caused by serogroup W reported before 2015. Prior to 2016, the proportion of cases of IMD caused by serogroup W ranged from 1–5% in the period 1997–2012, 8–10% in 2013–2014, to 21% of the total cases of IMD in 2015.

The number and proportion of cases of IMD caused by serogroup Y in 2016 (n=40, 17% of total IMD) was also the highest reported by the AMSP, almost double the number of IMD cases reported in the previous year (22 cases), and a four-fold increase in the average number of annual serogroup Y cases reported before 2015. Prior to 2016, the proportion of cases of IMD caused by serogroup Y ranged from 1–5% in the period 1997–2010, 6–11% in 2011–2014, and was 13% of the total cases of IMD in 2015.

Of the 107 cases of IMD cases caused by serogroup W in 2016, 47 cases (44%) were reported from Victoria, where serogroup W represented 62% (47/76) of cases, 23 cases (22%) were reported in New South Wales, representing 33% (23/69) cases and 14 cases (13%) were reported in Western Australia, representing 67% (14/21) cases. This is in contrast to previous years where, serogroup B has been predominant in these states. Serogroup W was reported in all jurisdictions, except in the Northern Territory in 2016.

Of the 40 cases of IMD caused by serogroup Y in 2016, 15 cases (38%) were reported from New South Wales, where this serogroup represented 22% (15/69 cases) of cases. Thirteen cases (33%) were reported in Queensland, representing 31%

Figure 2: Proportion of serogroups of laboratory-confirmed invasive meningococcal disease, Australia, by year



(13/42 cases) of cases. Serogroup Y was reported in all jurisdictions except in the Northern Territory, Australian Capital Territory, and South Australia. Serogroup B was reported in all jurisdictions except in Tasmania, and continued to be the predominant serogroup amongst IMD cases reported in South Australia.

In 2016, the predominant serogroup for children less than 5 years continued to be serogroup B, however the proportion was the lowest since 2000. (Table 3, Figure 3). In young adults aged 20–24 years, IMD caused by serogroup B was also the predominant serogroup, however this proportion was lower than in 2014 (83%), 2007–2010 and 2012 (72–88%), but similar to 2011, 2013, and 2015 (62–67%). The proportion of IMD caused by serogroup B in the all other age groups was less compared with previous years, due to the large increase in the number of IMD cases caused by serogroup W and Y in these age groups.

In 2016, there was an increase in the number of IMD caused by serogroup W across all age

groups except those aged 10–14 years. For those aged more than 45 years, IMD caused by serogroup W was the predominant serogroup (49/83 cases or 57%). This is in contrast with previous years, where serogroup Y was predominant for this age group. There was also a large increase in the number and proportion of IMD cases caused by serogroup W (16/35 cases, 46%) in those aged 25–44 years, compared with previous years, where serogroup B was predominant (68–87% since 2007).

Genotyping

In 2016, genotyping results were available for 93% (225/243) of IMD cases (Tables 5 and 6). The predominant *porA* genotype for IMD cases caused by serogroup B was P1.7-2,4 (25 cases, 33% of serogroup B that were typeable), which is similar to 2015 (Figure 4). The predominant *porA* genotype for serogroup Y IMD cases was P1.5-1,10-1 (34 cases, 85% of serogroup Y IMD cases that were typeable).

Figure 3: Number of serogroups B, Y and W cases of laboratory-confirmed invasive meningococcal disease, Australia, 2016, by age




Figure 4: Number of porA genotypes for serogroup B in laboratory-confirmed cases of invasive meningococcal disease Australia, 2016




Table 5: Laboratory-confirmed cases of invasive meningococcal disease, Australia, 2016, by *porA* genotype

2016 AMSP GenotypeporA	No. PER SEROGROUP					
	B	C	Y	W135	ND	TOTAL
P1.5,2	0	2	0	96	2	100
P1.5,2-59	0	0	0	1	0	1
P1.5,10-1	1	0	0	0	0	1
P1.5-1,2-2	0	0	2	0	0	2
P1.5-1,10-1	0	0	34	0	0	34
P1.5-1,10-4	0	0	0	2	0	2
P1.5-2,10-1	0	0	3	0	0	3
P1.5-2,10-2	1	0	0	0	0	1
P1.5-2,10-29	0	0	1	0	0	1
P1.5-8,10-4	1	0	0	0	0	1
P1.5-9,2	0	0	0	1	0	1
P1.5-11,2	0	0	0	1	0	1
P1.7,16-26	6	0	0	0	1	7
P1.7-2,?	1	0	0	0	0	1
P1.7-2,4	25	0	0	0	0	25
P1.7-2,4-31	1	0	0	0	0	1
P1.7-2,13	1	0	0	0	0	1
P1.7-2,16-26	1	0	0	0	0	1
P1.7-4,1	1	0	0	0	0	1
P1.12-1,13	1	1	0	0	0	2
P1.12-1,16	2	0	0	0	0	2
P1.12-1,16-65	1	0	0	0	0	1
P1.17,16-26	1	0	0	0	0	1
P1.18-1,2-34	1	0	0	0	0	1
P1.18-1,3	2	0	0	3	0	5
P1.18-1,34	6	0	0	0	0	6
P1.18-1,34-5	1	0	0	0	0	1
P1.19,13-1	1	0	0	0	0	1
P1.19,15	1	0	0	0	0	1
P1.19-1,15	1	0	0	0	0	1
P1.21-7,?	1	0	0	0	0	1
P1.22,9	4	0	0	0	0	4
P1.22,14	9	0	0	0	0	9
P1.22,14-6	1	0	0	0	0	1
P1.22,14-22	1	0	0	0	0	1
P1.22-1,14	1	0	0	0	0	1
P1.22-15,14	1	0	0	0	0	1
TOTAL	75	3	40	104	3	225

Table 6: Distribution of *porA* genotype laboratory-confirmed cases of invasive meningococcal disease, Australia, 2016, by state or territory

Genotype <i>porA</i>	No. PER SEROGROUP PER STATE							
	NSW	QLD	VIC	SA	WA	ACT	TAS	NT
P1.5,2	21W, 1C	12W	44W, 1C, 1NG	4W	14W	1W	1W	
P1.5,2-59			1W					
P1.5,10-1				1B				
P1.5-1,2-2		1Y		1Y				
P1.5-1,10-1	15Y	10Y	8Y		1Y			
P1.5-1,10-4		2W						
P1.5-2,10-1		2Y	1Y					
P1.5-2,10-2		1B						
P1.5-2,10-29							1Y	
P1.5-8,10-4			1B					
P1.5-9,2	1W							
P1.5-11,2			1W					
P1.7,16-26		3B	4B					
P1.7-2,?	1B							
P1.7-2,4	3B	3B	3B	15B	1B			
P1.7-2,4-31			1B					
P1.7-2,13					1B			
P1.7-2,16-26		1B						
P1.7-4,1					1B			
P1.12-1,13	1C							1B
P1.12-1,16	2B							
P1.12-1,16-65		1B						
P1.17,16-26	1B							
P1.18-1,2-34			1B					
P1.18-1,3			1B				3W	1B
P1.18-1,34	4B	1B			1B			
P1.18-1,34-5			1B					
P1.19,13-1				1B				
P1.19,15		1B						
P1.19-1,15	1B							
P1.21-7,?	1B							
P1.22,9	2B		1B	1B				
P1.22,14	4B	2B	3B					
P1.22,14-6		1B						
P1.22,14-22	1B							
P1.22-1,14	1B							
P1.22-15,14			1B					

There were 105 serogroup W IMD cases that were able to be genotyped, 97 of these (92%), had the *porA* antigen encoding gene type P1.5,2. Multilocus sequence typing showed that, of these 72% (70/97) were sequence type (ST)-11 — the same strain type as the hypervirulent serogroup W strain reported in the UK and South America since 2009^{4,5} (Table 7).

Antibiotic susceptibility testing

Antimicrobial susceptibility testing was able to be performed for 77% (189/243) of the IMD cases in 2016. Of these, 6% (11/189) were resistant to penicillin (MIC \geq 1 mg/L), the highest number and proportion of isolates with penicillin resistance reported by the AMSP. Only 4% of isolates tested were fully susceptible to penicillin (MIC \leq 0.03 mg/L), and 90% (170/189) of isolates were less sensitive to penicillin (MIC=0.06–0.5 mg/L). Of the isolates that were resistant to penicillin, all were serogroup W, and 9/11 were P1.5,2: ST-11, one was P.1.5,2: ST-12351, and one was P.1.5,2-59: ST-11. This represented 11% penicillin resistance in all serogroup W isolates available for testing (n=99). There was one isolate that had an elevated MIC to ceftriaxone (MIC=0.125 mg/L), the highest recorded by the AMSP. This isolate was W:P1.5,2, ST-11 and was less sensitive to penicillin. All isolates tested were susceptible to rifampicin and ciprofloxacin.

Discussion

In 2016, there were 243 cases of laboratory confirmed IMD, representing 96% of the number of notifications to the NNDSS.² The number of laboratory-confirmed IMD in 2016 represents a 40% increase cases compared with the previous year. The number of IMD cases has been steadily rising since the nadir in 2013, however the number of cases reported this year remains less than half of the peak number of IMD cases reported in Australia in 2002 (n=684). The introduction of the serogroup C vaccine to the national immunisation schedule in 2003 has resulted in a very large and sustained reduction in the number and proportion of serogroup C IMD cases in this country, and in 2015 the number and proportion of IMD cases caused

by serogroup C was the lowest ever reported by the AMSP. In early 2014, a recombinant multi-component meningococcal B vaccine became available in Australia.⁶ This vaccine is not on the national immunisation program but is available for purchase privately. Therefore uptake is elective and the impact of its introduction is yet to be determined in this country.

A number of changes in IMD epidemiology were observed in Australia in 2016. There was a notable increase in the number and proportion of IMD cases caused by serogroup W and serogroup Y, and a marked decrease in the number and proportion of cases caused by serogroup B. The incidence of serogroup W and serogroup Y was the highest ever reported by the AMSP.

In addition, as reported by the AMSP in 2015, the primary peak of IMD was observed in adults aged 45 years or older and was due to the increased number of IMD cases caused by serogroup W and serogroup Y in this age group. However these serogroups also increased in frequency in all age groups in 2016. Secondary disease peaks were observed in those aged less than 5 years, and in adolescents aged 15–19 years.

Serogroup W represented 44% of all laboratory-confirmed IMD cases in 2016, with the highest proportions in Victoria (44%), Queensland (33%), New South Wales (22%) and Western Australia (13%). Typing of these isolates found that the predominant circulating strain of serogroup W, had the *porA* antigen encoding gene type P1.5,2 and was sequence type (ST)-11. This is same genotype as the hypervirulent serogroup W strain that emerged in the United Kingdom and South America in 2009^{4,7} and has spread to now account for 25% of IMD in the UK in 2014/15, and 59% of all cases in Chile in 2012. This serogroup W strain is now considered endemic in these regions and is associated with atypical presentations, more severe clinical disease and a higher case fatality rate.⁷ The initial increase in serogroup W in these regions, as is now being demonstrated in Australia, was seen in older adults, but was subsequently reported in all age groups, particularly in adolescents and

Table 7: Laboratory-confirmed cases of serogroup W IMD, Australia, 2016, by sequence type (ST)

Sequence Type	W Genotype						Not typeable	Total
	P1.5,2	P1.5,2-59	P1.5-1,10-4	P1.5-9,2	P1.5-11,2	P1.18-1,3		
ST11	70	1	0	0	0	0	0	71
ST22	0	0	0	0	0	3	0	3
ST23	0	0	2	0	0	0	0	2
ST1287	7	0	0	0	0	0	0	7
ST3298	1	0	0	0	0	0	0	1
ST8345	1	0	0	0	0	0	0	1
ST8857	0	0	0	0	1	0	0	1
ST10651	0	0	0	1	0	0	0	1
ST12351	8	0	0	0	0	0	0	8
ST12590	1	0	0	0	0	0	0	1
Not typeable	9	0	0	0	0	0	2	11
Total	97	1	2	1	1	3	2	107

infants⁸. In response, vaccination programs have been implemented in both the United Kingdom and in Chile^{4,9}

An increase in IMD cases caused by serogroup Y was also observed in the eastern states of New South Wales, Queensland and Victoria in 2016. The predominant serogroup Y genotype (85%, 34/40) was P1.5-1,10-1, which increased in number and proportion compared with 2015 (70%, 14/20) and 2014 (50%, 6/12) whereas in previous years the serogroup Y genotype distribution was more heterogeneous. The emergence of serogroup Y and genotype Y:P1.5-1,10-1 has also been reported recently in Europe¹⁰. The phenotypic and genotypic characterization of the serogroup Y isolates is ongoing by the NNN.

Antimicrobial susceptibility testing of IMD isolates in 2016 showed an increase in penicillin resistance, the highest annual number and proportion recorded by the AMSP. The incidence of penicillin resistance in *N. meningitidis* in Australia has been less than 1% annually of IMD isolates tested in 1996-2014, rising to 3% in 2015, and 6% in 2016. The proportion of IMD isolates with penicillin MIC values in the less sensitive category has been increasing in recent years. These proportions ranged from 62–75% in

1996–2006; 67%–79% in 2007–2009; 78%–88% in 2010–2015, and in 2016, was 90%. In 2016, all isolates resistant to penicillin were serogroup W, representing 11% of all serogroup W isolates tested. Additionally, there was another serogroup W isolate with the highest ceftriaxone MIC recorded (0.125mg/L) by the AMSP. All IMD isolates were susceptible to rifampicin and ciprofloxacin.

The increase in IMD cases caused by serogroup W and serogroup Y, and the observed increase in antimicrobial resistance in serogroup W isolates are of significant concern. The NNN is continuing to lead further investigations with the Department of Health and the CDNA in to these observed changes and is closely monitoring the phenotypic and genotypic features of *N. meningitidis* causing IMD in Australia. Additional investigations including whole genome sequencing are in place to enhance IMD surveillance. The AMSP data are used for informing treatment guidelines and disease prevention strategies; and to monitor the effect of interventions.

Acknowledgements

Meningococcal isolates were received in the reference centres from many laboratories throughout Australia. The considerable time and effort involved in forwarding these isolates is recognised and these efforts are greatly appreciated. These data could not have been provided without this assistance and the help of clinical colleagues and public health personnel. The Australian Government Department of Health provided funding for the National Neisseria Network.

Members of the AMSP in 2016, to whom isolates and samples should be referred, and enquiries directed, are listed below.

Australian Capital Territory

P Collignon, S Bradbury
Microbiology Department
The Canberra Hospital
Yamba Drive
Garran ACT 2605
Telephone: +61 2 6244 2414
Email: peter.collignon@act.gov.au

New South Wales

MM Lahra, RP Enriquez, EA Limnios, TR Hogan, RL Kundu
Microbiology Department, New South Wales Health Pathology, The Prince of Wales Hospital
Barker Street, Randwick NSW 2031
Telephone: +61 2 9382 9084
Facsimile: +61 2 9382 9310
Email: monica.lahra@health.nsw.gov.au

M Maley, J Mercer, R Porritt
Department of Microbiology and Infectious Diseases
SSWPS
Locked Mail Bag 7090
Liverpool BC NSW 1871
Telephone: +61 8738 5124
Facsimile: +61 2 8738 5129
Email: Joanne.Mercer@sswahs.nsw.gov.au or
Robert.Porritt@sswahs.nsw.gov.au

Northern Territory

R Baird, K Freeman
Microbiology Laboratory
Territory Pathology
Royal Darwin Hospital
Tiwi NT 0810
Telephone: +61 8 8922 8167
Facsimile: +61 8 8922 7788
Email: rob.baird@nt.gov.au

Queensland

G Robertson, J Bates, H Smith, V Hicks
Public Health Microbiology
Forensic and Scientific Services
39 Kessels Road
Coopers Plains Qld 4108
Telephone: +61 7 3274 9101
Facsimile: +61 7 3274 9175
Email: john_bates@health.qld.gov.au

South Australia

I Bastian, A Lawrence
Microbiology and Infectious Diseases
SA Pathology
Frome Road, Adelaide, SA 5000
Telephone: +61 8 2223503
Facsimile: +61 8 2223543
Email: andrew.lawrence@health.sa.gov.au

Tasmania

L Cooley, B McEwan
Department of Microbiology and Infectious Diseases
Royal Hobart Hospital
48 Liverpool Street
Hobart Tasmania 7000
Telephone: +61 3 6222 8656
Email: belinda.mcewan@dhhs.tas.gov.au

Victoria

B Howden, K Stevens
Microbiological Diagnostic Unit Public Health Laboratory
Department of Microbiology and Immunology
The University of Melbourne

Parkville Victoria 3052
 Telephone: +61 3 8344 5701
 Facsimile: +61 3 8344 7833
 Email: kerries@unimelb.edu.au

Western Australia

AD Keil, J Bew
 Department of Microbiology
 Princess Margaret Hospital for Children
 1 Thomas Street
 Subiaco WA 6008
 Telephone: +61 8 9340 8273
 Facsimile: +61 8 9380 4474
 Email: tony.keil@health.wa.gov.au or jane.bew@health.wa.gov.au

Author details

Monica M Lahra^{1,2}

Rodney Enriquez¹

1. Neisseria Reference Laboratory and World Health Organisation Collaborating Centre for STD, Sydney. Department of Microbiology, South Eastern Area Laboratory Services, The Prince of Wales Hospital, Randwick, 2030, NSW Australia.

2. School of Medical Sciences, Faculty of Medicine, The University of New South Wales, NSW, 2052 Australia

Corresponding author: Professor Monica Lahra, Microbiology Department, SEALS, Director, Neisseria Reference Laboratory and WHO Collaborating Centre for STD, Level 4, Campus Centre, The Prince of Wales Hospital, RANDWICK NSW, 2031. Email: monica.lahra@health.nsw.gov.au

References

1. Meningococcal Isolate Surveillance Australia, 1994. [Internet]. National Neisseria Network. 1995.

2. NNDSS. Number of notifications of Meningococcal disease (invasive), received from

State and Territory health authorities in the period of 1991 to 2012 and year-to-date notifications for 2014.: National Notifiable Diseases Surveillance System; 2014.

3. Communicable Diseases Network Australia. Invasive Meningococcal Disease: CDNA National Guidelines for Public Health Units 2015. Available from: <http://www.health.gov.au/internet/main/publishing.nsf/content/cdna-song-imd.htm>.

4. Abad R, Lopez E, Debbag R, et al. Sero-group W meningococcal disease: global spread and current affect on the Southern Cone in Latin America. *Epidemiol Infect.* 2014;142:2461-70.

5. Ladhani S, Beebejuan K, Kucidarme H, et al. Increase in endemic *Neisseria meningitidis* capsular group W sequence type 11 complex associated with severe invasive diseases in England and Wales. *Clinical Infectious Diseases.* 2015;60:578-85.

6. Australian Government Department of Health. Meningococcal Disease. Immunise Australia Program. 2015. Available from: <http://www.health.gov.au/internet/immunise/publishing.nsf/Content/immunise-meningococcal>

7. Ladhani SN, Beebejaun K, Lucidarme J, Campbell H, Gray S, Kaczmariski E, et al. Increase in endemic *Neisseria meningitidis* capsular group W sequence type 11 complex associated with severe invasive disease in England and Wales. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America.* 2015;60(4):578-85.

8. Araya P, Fernandez J, Del Canto F, et al. *Neisseria meningitidis* ST-11 Clonal Complex Chile, 2012. *Emerging Infectious Diseases.* 2015;21(2):339-41.

9. Campbell H, Saliba V, Borrow R, Ramsay M, SN L. Targeted vaccination of teenagers following continued rapid endemic

expansion of a single meningococcal group W clone (sequence type 11 clonal complex), United Kingdom 2015. *Euro Surveill.* 2015;20(28):pii=21188.

10. Broker M, Jacobsson S, Kuusi M, Pace D, Simoes MJ, Skoczynska A, Taha M-K, et al. Meningococcal serogroup Y emergence in Europe: update 2011. (2164-554X (Electronic)). *Hum Vaccin Immunother.* 2012 Dec 1; 8(12): 1907–1911

Submit an Article

You are invited to submit your next communicable disease related article to the Communicable Diseases Intelligence (CDI) for consideration.

More information regarding CDI can be found at:
<http://health.gov.au/cdi>.

Further enquiries should be direct to:
cdi.editor@health.gov.au.